

Preliminary data, not for publication or official use

# Project Tycho

*Preliminary data for the state of Texas*

## Tycho database beta test version

*The data presented in this report are of preliminary nature and should not be used for publication or other types of official use*

June 2011

University of Pittsburgh Graduate School of Public Health



University of Pittsburgh



## Table of Content

Acknowledgements	page 3
Contact information	page 4
Introduction	page 5
Vision and goal	page 7
Collaboration	page 8
Summary of methods	page 9
Overview of data available for the state of Texas	page 12
Disease specific data for Texas	page 19
- AIDS	page 19
- Chickenpox	page 20
- Chlamydia	page 22
- Coccidioidomycosis	page 23
- Cryptosporidiosis	page 24
- Diphtheria	page 25
- Escherichia Coli	page 28
- Gonorrhea	page 29
- Haemophilus Influenzae	page 30
- Hepatitis	page 31
- Influenza	page 33
- Legionellosis	page 34
- Listeriosis	page 35
- Lyme disease	page 36
- Malaria	page 37
- Measles	page 38
- Meningitis	page 40
- Meningococcal disease	page 42
- Mumps	page 44
- Pneumonia	page 46
- Poliomyelitis	page 48
- Rabies in animals	page 50
- Rubella	page 52
- Salmonellosis	page 53
- Scarlet fever	page 54
- Shigellosis	page 57
- Streptococcal disease, invasive group A	page 58
- Streptococcal sore throat	page 59
- Streptococcus pneumoniae, invasive disease	page 60

# Preliminary data, not for publication or official use

## Disease specific data for Texas, continued

- Syphilis	page 61
- Tetanus	page 62
- Tularemia	page 63
- Tuberculosis	page 64
- Typhus fever	page 65
- Typhoid fever	page 68
- West nile disease	page 69
- Whooping cough	page 70
<b>Diseases with fragmented data</b>	<b>page 72</b>

## Acknowledgements

The development of this project would not have been possible without data collected every week by staff from city-, county-, state- and federal public health departments. We would like to acknowledge their invaluable contribution to the overall body of surveillance data that were digitized during this project. Particularly, we acknowledge staff from the Council of State and Territorial Epidemiologists (CSTE) and the US Centers for Disease Control for their overall efforts in disease surveillance and their particular input in this project.

We also acknowledge the help of many contributors that have contributed to the digitization and processing of weekly surveillance data included in this database. Digital Divide Data: Linda Thomas, Eric Gold, Heak Hok, Villa Kem and Hetel Patel. University of Pittsburgh Graduate School of Public Health: Steve Wisniewsky, Heather Eng, Sharon Crow, David Galloway, Jia Xu and Raaka Kumbhakar. University of Pittsburgh School of Information Sciences: Vladimir Zadorozhny, Ying-Feng Hsu, David Paul Koenig, Fatimah Ridwan and Divyasheel Sharma. Johns Hopkins Bloomberg School of Public Health: Derek Cummings. Pittsburgh Supercomputing Center: David Graham and Andrew Adams.

This project was made possible with funding from the Bill & Melinda Gates Foundation through the Vaccine Modeling Initiative (grant ID 49276).

The Tycho development team at the University of Pittsburgh Graduate School of Public Health:

Willem G. van Panhuis  
John Grefenstette  
Shawn T. Brown  
Su Yon Jung  
Nian Shong Chok  
Anne Cross  
Donald S. Burke

## Contact information

The Tycho database is currently being tested and a login account will be required to use the database and website ([www.tycho.pitt.edu](http://www.tycho.pitt.edu)). All State Epidemiologists, CDC partners, other federal agencies and our research collaborators will be provided with user accounts. User accounts can also be provided upon specific request. Note that the quality of the data during the testing phase cannot be guaranteed to be sufficient for publication or official use.

**Point of contact for the Tycho project:**

Dr. Willem G. van Panhuis, MD PhD

Graduate School of Public Health

University of Pittsburgh

130 DeSoto Street

704 Parran Hall

Pittsburgh PA, 15261

Email: [wav10@pitt.edu](mailto:wav10@pitt.edu)

Tel: 412-624-7693

## Introduction

**Project Tycho** started in 2009 as part of the Vaccine Modeling Initiative (VMI) at the University of Pittsburgh Graduate School of Public Health, funded by the Bill & Melinda Gates foundation. This project aims to contribute to the availability of public health data for research and policy making. The vision for this project as described in the next section is a central access point for public health data of a detailed spatial and temporal scale for all countries. Currently, the Tycho database contains 10 million records that each represent a weekly report from a location for a specific disease. These records were extracted from weekly US Nationally Notifiable Disease Surveillance Reports between 1888 and 2009 (6300 reports) using double data entry (200 million keystrokes). The current database includes a total of 90 million reported cases and 4 million reported deaths due to notifiable diseases in the United States for the last 122 years.

Project Tycho is named after the Danish nobleman Tycho Brahe (1546 – 1601), who meticulously collected astronomical data. After Brahe's death, his assistant Kepler used these data to develop area laws of planetary motion.

## Background

***"No health department, State or local, can effectively prevent or control diseases without knowledge of when, where and under what conditions cases are occurring"***

(Quote stated above weekly surveillance reports published in the Public Health Reports, 1888-1951)

This quote has accompanied weekly reports on the prevalence of notifiable diseases in the United States as long as these were published in the Public Health Reports, emphasizing the importance of data for disease control policy making. In the current day and age, this statement has lost none of its significance. On the contrary, as infectious diseases continue to pose challenges to the global public health system in an increasingly interconnected global society. The public health system has responded to this challenge by evermore sophisticated disease surveillance systems. The availability of internet and information technology has facilitated collection of detailed data on infectious disease incidence, the spread of pathogens, disease determinants, health behavior, etc. New technologies such as remote sensing and mobile reporting systems have advanced measurement of these factors in both high and low resource settings. The enormous wealth of public health information that is currently being accrued also poses its own challenges for data storage, management, preservation and dissemination. Public health agencies will need to allocate resources and develop capacity to specifically address these issues.

Although the importance of public health data collection has been widely recognized, the dissemination of data has been neglected for many reasons. The above quote does not only imply data collection but also its use and dissemination as knowledge on disease occurrence can only be derived from data after processing and analysis. Not only has disease surveillance become more sophisticated, analytical tools have changed as well and now include computational models to assess disease transmission and to evaluate disease control options. These methods have introduced the opportunity for data intensive studies of fine spatial-temporal patterns using dynamic models. Disease surveillance data at such scale are often not available for analysis for multiple reasons. First, surveillance data are not collected for research purposes but for disease monitoring and planning. Secondly, public health staff often lack time or capacity to prepare these data for analysis. Third, surveillance methodology often lacks transparency and standardization, reducing options for scientific analyses. Fourth, disease surveillance data are often

# Preliminary data, not for publication or official use

kept confidential and ownership or data use requirements are often unclear. Finally, archiving and preservation of surveillance data are often not a priority and disaggregated records may get lost, losing the opportunity to ever use these data for analysis.

The Tycho project aims at overcoming some of these challenges by providing open access to large public health datasets of great spatial-temporal detail. Open access to these data will provide great opportunities for scientific analysis and better informed policy making on disease control.

## **The Tycho database and website**

As described in the methods section, a large heterogeneity in reporting practice was observed over the past 122 years of weekly disease surveillance and substantial efforts were made to clean and standardize these data. The current online database includes all cleaned and standardized data in a format that allows comparisons across time and space. It does not include parts of the data that require additional cleaning and research such as classification of disease subcategories.

Each record in the database provides a reported number of cases or deaths due to a specific disease for a specific location, for a certain week. The database can be searched online ([www.tycho.pitt.edu](http://www.tycho.pitt.edu)) by selecting a dataset, an aggregation method and an output format. Data can be viewed in tables, graphics and maps and these can be downloaded for further use.

## **Data availability**

The data provided by the Tycho database reflect the weekly US Nationally Notifiable Disease Surveillance System between 1888 and 2009. It includes all diseases that were reported as part of this system, but it does not yet include records from other surveillance systems such as the annual system or disease specific subsystem (such as the influenza surveillance system).

The available data also reflects changes in reporting practices over time. This explains why city or state level reports are only available for a certain time period (1888-1953 for cities and 1927-2009 for states) and why morbidity and mortality reports are available for different time periods. We are currently working on inclusion of additional data as described in the section on collaboration.

## **Testing and release of the Tycho database**

The Tycho database is currently in a beta testing phase that will start during the 2011 annual meeting of the Council of State and Territorial Epidemiologists (CSTE) in Pittsburgh June 12-16. This phase will end with a release to the general public in the fall of 2011. During the testing phase, invited users will be provided with login accounts that will enable full use of all features of the website and database. Invited users will include all state epidemiologists, partners in the CDC and other federal agencies as well as research collaborators. During this phase, data cleaning and standardization will continue. The quality of the data will not be sufficient for publication or official use during this phase.

## **Future developments**

After optimization of the database and website during the testing phase, the first version of the Tycho database will be released to the general public. After that, efforts to clean and standardize the data will continue and more data will be released in subsequent versions. In addition, ongoing and new collaborations for inclusion of new data will allow gradual expansion of the dataset over time and increased opportunities for data driven analysis and policy making.

## Vision and goal

The vision for project Tycho is based on the value of open access to detailed, disaggregated public health data for scientific analysis and policy making. Starting with 122 years of weekly US surveillance data, it is our hope that this example will convince public health authorities worldwide of the value of this resource and will stimulate similar contributions from other countries.

In the past, data sharing and archiving have not been a priority in public health or among health sciences in general. Continuous and diligent public health data collection has led to a large pool of disaggregated data scattered over the world including the internet, ranging from paper archives to basements of local health departments. The need for open access to a central repository of public health data has been recognized and both benefits and challenges of open access to data in general have been well described. Open access to data could lead to increased accountability, transparency, innovation, collaboration, cost-effectiveness, replication of results, the development of new methods and insights, and a narrower data access gap between low- and higher income countries. Ultimately, this would result in the advancement of science and technology to the benefit of all.

Challenges to open access include inconsistent formats, lack of annotation or metadata, lack of information technology (IT) capacity, a lack of incentive, and cultural norms around data ownership. Open access to public health data in particular is confronted by decentralized archiving of disease reports, lack of time and capacity for data management, degraded physical format of records and archives, and governmental level legal constraints. Multiple principles and requirements for a central open access repository for public health data have been described and include (1) capacity building for data management in low- and middle income countries, (2) feedback loops to data contributors, (3) sustainability, (4) common standards, (5) interoperability, and (6) user-friendliness.

The ultimate goal of the Tycho project is to provide a central global public health data access point. Historical as well as current public health data are of great value if archived and accessible for research and analysis. Open access will enable the use of analytical capacity from around the globe which will lead to new discoveries of disease patterns and control policies. Furthermore, a central data access point will facilitate data archiving and preservation into the future which will be an increasing need in a data rich public health environment.

## Collaboration

The application of the surveillance data from the Tycho database as well as the inclusion of new data will require a wide range of collaborations with public health offices in the US and abroad. The Tycho project has currently made the first step of digitizing 122 years of weekly US surveillance reports that had already been published. The next step will be the completion of the existing dataset as well as the inclusion of new data from unpublished sources.

This report provides an overview of data available from the Tycho database for the state of Texas. This includes all data that has ever been published at state or city level for Texas in the weekly US Nationally Notifiable Disease Surveillance System. As you will find, the data availability will vary greatly over time, between city and state level reports, between diseases and between morbidity vs. mortality reports.

The Tycho development team would look forward to work with state health departments in the US to:

1. Better understand the current Tycho data for each state (eg. by collecting historic documentation)
2. Provide better usability and applications of current Tycho data for each state by continued joint development of website and database features.
3. Collect and/or digitize official, confirmed data to validate the current (preliminary) weekly data.
4. For each state, collect and/or digitize new data that has not yet been included. For example all city level reports discontinued in 1953 and completion of city level data until 2009 would greatly increase opportunities for analysis of disease patterns and trends.
5. Provide support to state health offices to manage, preserve and provide access to public health data.

## Summary of methods

Detailed documentation on the methods used for the Tycho database has been provided on the website ([www.tycho.pitt.edu](http://www.tycho.pitt.edu)). This section describes these methods in short.

### Data collection

Weekly reports that contain tables on the occurrence of nationally notifiable diseases have been published since 1888 by public health authorities at the Federal level in various journals. Table 1 provides the list of publications and the responsible agency since 1888. All weekly nationally notifiable disease reports between 1888 and 1951 could be retrieved from the PubMed Central repository of the National Library of Medicine <sup>1</sup>. For 1995 to 2009, these reports could be retrieved from the MMWR digital archive on the CDC website <sup>2</sup>. Most weekly reports between 1952 and 1995 could be retrieved from the HathiTrust Digital Library <sup>3 4</sup>, but many could not be found and had to be copied from hard copies of MMWR issues in the University of Pittsburgh library.

Table 1, publications and responsible Federal agencies for nationally notifiable disease reports

Time period	Publication title	Responsible federal agency
1888-1889	Weekly Abstract of Sanitary Reports	US Marine Hospital Service
1890-1895	Abstract of Sanitary Reports	US Marine Hospital Service
1896-1901	Public Health Reports	US Marine Hospital Service
1902-1911	Public Health Reports	US Public Health and Marine Hospital Service
1912-1951	Public Health Reports	US Public Health Service
1952-1960	Morbidity and Mortality Weekly Report	National Office of Vital Statistics, US Public Health Service
1961-1969	Morbidity and Mortality Weekly Report	Communicable Disease Center
1970-1991	Morbidity and Mortality Weekly Report	Center for Disease Control
1992-2009	Morbidity and Mortality Weekly Report	Centers for Disease Control and Prevention

### Inclusion criteria

Weekly reports of each year were reviewed systematically to assess the diseases reported. We included all tables that provided disease specific information by week for US cities, townships, counties or states. Tables that provided summary or aggregated information by month, year or at the national level were not included. Similarly, tables that did not contain disease specific information (such as all cause mortality) were not included.

### Data entry

Weekly reports were downloaded or scanned as PDF files and selected tables with notifiable disease reports were entered into computer spreadsheets in a highly standardized fashion using double data entry. During the second round of data entry, operators could not see what had been entered in the first round and could not continue if the system detected a discrepancy between the second and first entry

<sup>1</sup> <http://www.ncbi.nlm.nih.gov/pmc/tocrender.fcgi?journal=333&action=archive>

<sup>2</sup> <http://www.cdc.gov/mmwr>

<sup>3</sup> <http://catalog.hathitrust.org/Record/003910026>

<sup>4</sup> <http://catalog.hathitrust.org/Record/003843660>

# Preliminary data, not for publication or official use

for a specific value. Such discrepancies could only be resolved by checking the PDF file and try again or discussion with the group leader and verification of a value in the source documents.

## **Quality control for data entry**

The accuracy of data entry was checked at various levels. First, completeness of data was verified by comparing the content of entered data with PDF sources files. Secondly, accuracy of data entry was verified by multiple rounds of comparing random samples of entered files with PDF source files. Thirdly, data formatting was verified by various checks to ensure appropriate formatting for data loading.

## **Data loading and standardization**

All data was entered in Excel spreadsheets and various components of these spreadsheets were loaded in data files. Table titles, column headers, place names and reported numbers were loaded in separate files. These files were used to extract information on each reported number, including:

1. the disease reported
2. the disease subcategory reported
3. cases or deaths reported
4. the reporting location (name, state and type of location)
5. the time period for which a number was reported
6. the date of publication of the original weekly report associated with a reported number

## **Integration**

All reported numbers and extracted information was integrated in one database with one record per reported number and associated information.

## **Post-processing quality control**

After integration of all data in one database, checks were performed to detect duplicate reports and data inconsistencies. Duplicate records were removed and inconsistencies resolved by verification with original PDF source files.


## **Data filtering**

The digitized version of all historical weekly US nationally notifiable disease surveillance records is a very heterogeneous dataset. It took a substantial number of data processing protocols to standardize time and space variables as well as disease names. All standardized records have been separated from non-standard records by a filter. All remaining heterogeneity in non-standard records is inherent to the surveillance system and can only be standardized after further analysis will have been completed (eg. remaining heterogeneity in reporting periods for reports before 1953 and standardization of disease subcategories). The largest proportion of data (>4 million records) has been standardized however and has been made available in the current testing version.

## **Data visualization**

Figures in this report were made with the R system, version 2.9.2 and the maps were generated by the GAIA platform developed at the University of Pittsburgh Graduate School of Public Health in collaboration with the Pittsburgh Supercomputing Center (PSC). See <http://midas-pitt.psc.edu/gaia> for more information.

# Preliminary data, not for publication or official use



**Figure 1, Schematic of the Tycho data architecture.** Access to a consistent, standardized subset of the digitized US weekly surveillance reports will be provided for beta testing through a website that will allow querying and downloading of data. Data from the development database will be pushed through a filter to feed the Tycho database.

The current Tycho database will provide the reported number of cases or deaths reported by city or state health authorities to the federal health agency for all weeks between 1888 and 2009. Currently, only standardized, consistent data are being released for testing, according to the following criteria:

1. Only weekly reports are provided (reports for other periods such as 10 days, 2 weeks, 1 month, 1 year) are currently not included due to lack of comparability with the weekly reports.
2. Only reports that were published in the same year as the reporting period were included (this excludes updates or comparisons that were published more than a year after the original reporting period)
3. Only numbers were included for which information about the location, time period and disease could be extracted.
4. Only numbers or time periods for which no disease subcategories were reported were included.

Each of the steps described above have been described in detailed protocols that are posted in the documentation part of the Tycho website ([www.tycho.pitt.edu](http://www.tycho.pitt.edu)).

## Website

The Tycho website features simple and advanced searches of the data, visualizations and downloading of machine readable files. A dataset can be selected by specifying: 1) the disease of interest, 2) the location(s) of interest and 3) the time period. In a second step, the type of aggregation and length of increments can be specified and in a third step, the type of output (table, graphic or map) can be selected. After output has been generated, data can be downloaded.

As mentioned above, data availability varies largely by disease, time period and location. We aimed to restrict selection options by data availability to avoid "no results available."

Access to the most of the website will be password protected during the testing phase. For invited users after login, full functionality of the database and website will be available. During this phase however, the data quality will not be sufficient for publication or official use.

## Overview of data available for the state of Texas

In this section, an overview of data available for the state of Texas will be provided. As described earlier, data availability depends entirely on historical reporting practices in the weekly US Nationally Notifiable Disease Surveillance System between 1888 and 2009. For this report, we only included morbidity reports (cases) to improve the format and limit the size of this report. As mentioned earlier, the analysis and standardization of has not been completed yet for all records in the database and preliminary data are provided here that may not yet be available in the online database.

This section will provide a general overview of data availability. The next sections will provide disease specific data from state and city level reports of Texas. Data for a maximum of 3 major cities were provided depending on availability (city data was only reported until 1953). Some summary data will be provided at the end of this report, for a subset of diseases for which data was fragmented over time.

Table 1 lists the number of weekly state or city reports that are available for Texas per each disease and subcategory.

**Table 1, Number of weekly state or city reports per disease and subcategory**

Disease	City	State
Aids	-	941
Anthrax	-	142
Botulism	-	1
Brucellosis [undulant fever]	1	816
Chickenpox [varicella]	454	792
Chlamydia	-	625
Coccidioidomycosis	-	11
Cryptosporidiosis	-	423
Dengue	9	-
Diphtheria	2093	2314
Dysentery		
<i>Amebic</i>	-	333
<i>Bacillary</i>	-	333
<i>Unspecified</i>	-	325
Encephalitis		
<i>Lethargic</i>	405	-
<i>Post infectious</i>	-	503
<i>Primary [infectious] including unspecified</i>	352	1607
Escherichia coli		
<i>EHEC 0157</i>	-	125
<i>EHEC non-0157</i>	-	86
<i>EHEC non serogrouped</i>	-	100
<i>O157:H7 NETSS</i>	-	240
<i>O157:H7 PHLIS</i>	-	159
<i>STEC</i>	-	175

# Preliminary data, not for publication or official use

**Table 1.** Number of weekly state or city reports per disease and subcategory, *continued*

Disease	City	State
Giardiasis	-	13
Gonorrhea		
<i>Civilian</i>	-	755
<i>Unspecified</i>	-	1007
Haemophilus influenzae		
<i>Age &lt;5 non-serotype B</i>	-	16
<i>Age &lt;5 serotype B</i>	-	81
<i>All ages all serotypes</i>	-	433
Hepatitis		
<i>Acute type A</i>	-	356
<i>Acute type B</i>	-	358
<i>Acute type C</i>	-	112
<i>Acute type NA NB [including C]</i>	-	14
<i>All types, &lt;20 years</i>	-	206
<i>All types, &gt;=20 years</i>	-	102
<i>All types, all ages</i>	25	694
<i>Type A [infectious]</i>	-	1670
<i>Type B [serum]</i>	-	1484
<i>Type NA NB [including C]</i>	-	752
<i>Type unspecified</i>	-	904
Influenza	961	1185
Legionellosis	-	733
Leprosy	1	509
Listeriosis	-	182
Lyme disease	-	690
Malaria		
<i>Civilian</i>	-	98
<i>Unspecified</i>	-	1723
Measles		
<i>Imported</i>	-	562
<i>Indigenous</i>	-	629
<i>Unspecified</i>	1878	2956
Meningitis		
<i>Aseptic</i>	-	1696
<i>Meningococcus</i>	864	1266
<i>Unspecified</i>	290	5

# Preliminary data, not for publication or official use

**Table 1.** Number of weekly state or city reports per disease and subcategory, *continued*

Disease	City	State
Meningococcal disease		
<i>All serogroups</i>	-	42
<i>Invasive all serogroups</i>	-	182
<i>Invasive serogroup unknown</i>	-	39
<i>Serogroup A C Y and W-135</i>	-	2
<i>Serogroup B</i>	-	2
<i>Serogroup unspecified</i>	72	2420
Mumps	454	1684
Pellagra	406	-
Pneumonia		
<i>Lobar</i>	27	-
<i>Unspecified</i>	202	169
Poliomyelitis		
<i>Non paralytic</i>	-	347
<i>Paralytic</i>	-	821
<i>Total</i>	1175	1968
Psittacosis	-	22
Rabies in animals	95	2484
Rocky mountain spotted fever	2	1407
Rubella	-	1503
Salmonellosis		
<i>NETSS</i>	-	119
<i>PHLIS</i>	-	128
<i>Unspecified</i>	-	359
Scarlet fever		
<i>Including streptococcal sore throat</i>	96	559
<i>Unspecified</i>	1922	1277
Shigellosis		
<i>NETSS</i>	-	121
<i>PHLIS</i>	-	124
<i>Unspecified</i>	-	378
Streptococcal disease, invasive group a	-	365
Streptococcal sore throat	-	102
Streptococcus pneumoniae invasive disease		
<i>Drug resistant &lt;5 years</i>	-	28
<i>Non drug resistant &lt;5 years</i>	-	201
Syphilis		
<i>Civilian primary and secondary</i>	-	738
<i>Congenital</i>	-	173
<i>Primary and secondary</i>	-	1056
Tetanus	-	731

# Preliminary data, not for publication or official use


**Table 1.** Number of weekly state or city reports per disease and subcategory, *continued*

Disease	City	State
Toxic shock syndrome	-	186
Tuberculosis [phthisis pulmonalis]		
<i>New active</i>	-	140
<i>Unspecified</i>	556	1309
Tularemia	1	1419
Typhoid fever [enteric fever]		
<i>Including paratyphoid fever</i>	339	709
<i>Unspecified</i>	1662	2519
Typhus fever		
<i>Endemic</i>	10	370
<i>Unspecified</i>	34	204
West nile disease		
<i>Neuroinvasive</i>	-	170
<i>Non-neuroinvasive</i>	-	108
Whooping cough [pertussis]	1367	2251
Yellow fever	1	-

The diseases included in the weekly US Nationally Notifiable Disease Surveillance System varied largely over time and reflected the historical social-political priorities of each time period. Note that the diseases in the weekly system were a subset of all diseases included in the annual Notifiable Disease Surveillance System (that was not entered as part of this project).

The number of locations reporting diseases also varied over time depending on the type of reports submitted. Figure 1 on the next page provides an overview of the number of unique locations reporting each disease per week between 1888 and 2009. For the remainder of this report, the state of Texas was considered a unique reporting location in addition to the individual city reports. Figure 1 indicates that many locations (mostly cities) were included in the reporting system before 1925, after which the number of cities dropped substantially. After 1953, only state reports were included and city reports were discontinued. These patterns were observed nationwide.

# Preliminary data, not for publication or official use




**Figure 1,** The number of locations (in color code) reporting on each disease per week between 1888 and 2009

For the majority of the time period between 1888 and 2009, only state level reports were available. It would be of great interest to complete the collection of city data after 1953 and we will seek collaboration with state health offices for this.

The map in figure 2 (next page) illustrates the geographical distribution of the available city data. It displays the number of weekly reports for any disease available per location that was included (and for which coordinates could be derived)


Preliminary data, not for publication or official use



**Figure 2,** This map displays the number of weekly reports available for any disease per city (note that city reports were only available until 1953)

# Preliminary data, not for publication or official use


Figure 3 below provides an overview of the number of diseases for which a weekly report was included at the state level and for individual cities. All cities for which at least 100 weekly reports were available have been listed. For most states, it shows a major drop in the number of cities after 1925 and after 1953, city reports were discontinued except for New York City, which became its own reporting jurisdiction.



**Figure 3,** The number of diseases (in color code) that were reported for the state and city level (state on top row) for each week between 1888 and 2009

## Disease specific data for Texas


### AIDS



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1, Summary information for AIDS**

Indicator	Texas
Report period	1984-2005
Total weeks	941
Total cases	60,126
Max. cases per year	6,436
Year (max)	1993
Max. cases per week	2,757
Week (max)	1993, wk 15
Average cases per year	2,733
95%CI	(2,064-3,402)
Average cases per week	64
95%CI	(54-74)



**Figure D2,** Number of cases reported for AIDS per epidemiological week

## Chickenpox




**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1,** Summary information for Chickenpox

Indicator	Galveston	San antonio	Houston
Report period	1924-1932	1924-1932	1924-1932
Total weeks	453	437	404
Total cases	80	275	584
Max. cases per year	29	66	121
Year (max)	1925	1931	1927
Max. cases per week	6	10	51
Week (max)	1925, wk 08	1931, wk 09	1927, wk 05
Average cases per year			
<i>before 1995</i>	9	31	65
95%CI	(2-16)	(18-44)	(38-92)
<i>after 1995</i>	-	-	-
95%CI	-	-	-
Average cases per week			
<i>before 1995</i>	0	1	1
95%CI	(0-0)	(1-1)	(1-1)
<i>after 1995</i>	-	-	-
95%CI	-	-	-

# Preliminary data, not for publication or official use



**Figure D2,** Number of cases reported for Chickenpox per epidemiological week


## Chlamydia



**Figure D1**, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1, Summary information for Chlamydia**

Indicator	Texas
Report period	1996-2009
Total weeks	621
Total cases	710,074
Max. cases per year	79,668
Year (max)	2007
Max. cases per week	7,353
Week (max)	1999, wk 26
Average cases per year	50,720
95%CI	(38,505-62,935)
Average cases per week	1,143
95%CI	(1,045-1,241)



**Figure D2**, Number of cases reported for Chlamydia per epidemiological week

## Coccidioidomycosis




**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1,** Summary information for Coccidioidomycosis

Indicator	Texas
Report period	2003-2003
Total weeks	11
Total cases	3
Max. cases per year	3
Year (max)	2003


## Cryptosporidiosis



**Figure D1**, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1**, Summary information for Cryptosporidiosis

Indicator	Texas
Report period	1999-2009
Total weeks	423
Total cases	2,275
Max. cases per year	1,281
Year (max)	2008
Max. cases per week	454
Week (max)	2008, wk 45
Average cases per year	207
95%CI	(-36-450)
Average cases per week	5
95%CI	(2-8)



**Figure D2**, Number of cases reported for Cryptosporidiosis per epidemiological week

## Diphtheria



**Figure D1**, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1**, Summary information for Diphtheria

Indicator	Texas	Galveston	San antonio	Houston
Report period	1927-1980	1907-1953	1907-1953	1906-1953
Total weeks	2,314	1,611	1,475	1,557
Total cases	59,448	1,485	3,036	6,728
Max. cases per year	5,906	103	324	460
Year (max)	1933	1915	1927	1933
Max. cases per week	4,496	11	32	38
Week (max)	1970, wk 44	1921, wk 47	1919, wk 48	1929, wk 45
Average cases per year				
before 1940	2,594	45	97	189
95%CI	(1,815-3,373)	(37-53)	(68-126)	(133-245)
after 1940	609	10	41	69
95%CI	(290-928)	(4-16)	(22-60)	(43-95)
Average cases per week				
before 1940	54	1	2	5
95%CI	(50-58)	(1-1)	(2-2)	(5-5)
after 1940	14	0	1	2
95%CI	(9-19)	(0-0)	(1-1)	(2-2)






Figure D2, Number of cases reported for Diphtheria per epidemiological week



**Figure D3,** Total number of all cases reported for Diphtheria by each city for the entire time period


## Escherichia Coli



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Escherichia Coli (0157:H7 PHLIS, EHEC 0157, and STEC)

Indicator	Texas
Report period	1996-2009
Total weeks	459
Total cases	814
Max. cases per year	116
Year (max)	2000
Max. cases per week	24
Week (max)	2008, wk 19
Average cases per year	63
95%CI	(37-89)
Average cases per week	2
95%CI	(2-2)



**Figure D2,** Number of cases reported for Escherichia Coli per epidemiological week


## Gonorrhea



**Figure D1**, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1**, Summary information for Gonorrhea (Unspecified and Civilian)

Indicator	Texas
Report period	1972-2009
Total weeks	1,761
Total cases	1,608,599
Max. cases per year	86,255
Year (max)	1978
Max. cases per week	9,547
Week (max)	1996, wk 32
Average cases per year	42,332
95%CI	(34,320-50,344)
Average cases per week	913
95%CI	(879-947)



**Figure D2**, Number of cases reported for Gonorrhea per epidemiological week


## Haemophilus Influenzae



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Haemophilus Influenzae (All ages all serotypes)

Indicator	Texas
Report period	1996-2009
Total weeks	433
Total cases	105
Max. cases per year	47
Year (max)	2006
Max. cases per week	47
Week (max)	2006, wk 35
Average cases per year	8
95%CI	(1-15)
Average cases per week	0
95%CI	(0-0)



**Figure D2,** Number of cases reported for Haemophilus Influenzae per epidemiological week


## Hepatitis



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Hepatitis (Type A [Infectious], Acute type A, Type B [Serum], and Acute type B)

Indicator	Texas
Report period	1966-2009
Total weeks	2,060
Total cases	69,438
Max. cases per year	4,301
Year (max)	1970
Max. cases per week	2,696
Week (max)	1970, wk 44
Average cases per year	
<i>before 1990</i>	1,919
<i>95%CI</i>	(1,631-2,207)
<i>after 1990</i>	1,130
<i>95%CI</i>	(777-1,483)
Average cases per week	
<i>before 1990</i>	41
<i>95%CI</i>	(36-46)
<i>after 1990</i>	24
<i>95%CI</i>	(21-27)



**Figure D2,** Number of cases reported for Hepatitis per epidemiological week


## Influenza



**Figure D1**, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1, Summary information for Influenza**

Indicator	Texas
Report period	1927-1950
Total weeks	1,165
Total cases	818,588
Max. cases per year	83,899
Year (max)	1950
Max. cases per week	9,392
Week (max)	1943, wk 51
Average cases per year	34,108
95%CI	(23,440-44,776)
Average cases per week	703
95%CI	(636-770)



**Figure D2**, Number of cases reported for Influenza per epidemiological week


## Legionellosis



**Figure D1**, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1**, Summary information for Legionellosis

Indicator	Texas
Report period	1983-2009
Total weeks	733
Total cases	706
Max. cases per year	111
Year (max)	2004
Max. cases per week	42
Week (max)	2004, wk 34
Average cases per year	26
95%CI	(15-37)
Average cases per week	1
95%CI	(1-1)



**Figure D2**, Number of cases reported for Legionellosis per epidemiological week


## Listeriosis



**Figure D1**, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1**, Summary information for Listeriosis

Indicator	Texas
Report period	2000-2005
Total weeks	182
Total cases	156
Max. cases per year	65
Year (max)	2003
Max. cases per week	11
Week (max)	2001, wk 40
Average cases per year	26
95%CI	(1-51)
Average cases per week	1
95%CI	(1-1)



**Figure D2**, Number of cases reported for Listeriosis per epidemiological week


## Lyme Disease



**Figure D1**, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1**, Summary information for Lyme Disease

Indicator	Texas
Report period	1991-2009
Total weeks	690
Total cases	939
Max. cases per year	142
Year (max)	2004
Max. cases per week	72
Week (max)	2001, wk 40
Average cases per year	49
95%CI	(32-66)
Average cases per week	1
95%CI	(1-1)



**Figure D2**, Number of cases reported for Lyme Disease per epidemiological week


## Malaria



**Figure D1**, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1**, Summary information for Malaria (Unspecified)

Indicator	Texas
Report period	1952-2009
Total weeks	1,723
Total cases	3,728
Max. cases per year	818
Year (max)	1970
Max. cases per week	376
Week (max)	1970, wk 45
Average cases per year	81
95%CI	(39-123)
Average cases per week	2
95%CI	(2-2)



**Figure D2**, Number of cases reported for Malaria per epidemiological week

## Measles




**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1,** Summary information for Measles (Unspecified)

Indicator	Texas	Galveston	San antonio	Houston
Report period	1927-2002	1909-1953	1907-1953	1907-1953
Total weeks	2,956	1,229	1,414	1,312
Total cases	1,188,364	1,958	7,254	7,422
Max. cases per year	82,299	754	897	1,012
Year (max)	1958	1953	1953	1924
Max. cases per week	6,795	250	160	183
Week (max)	1958, wk 18	1944, wk 15	1953, wk 24	1924, wk 09
Average cases per year				
before 1970	26,623	63	202	212
95%CI	(19,677-33,569)	(10-116)	(123-281)	(119-305)
after 1970	847	-	-	-
95%CI	(-89-1,783)	-	-	-
Average cases per week				
before 1970	543	2	5	6
95%CI	(504-582)	(1-3)	(4-6)	(5-7)
after 1970	21	-	-	-
95%CI	(16-26)	-	-	-

Preliminary data, not for publication or official use



**Figure D2,** Number of cases reported for Measles per epidemiological week


## Meningitis



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Meningitis (Meningococcus)

Indicator	Texas	Houston
Report period	1927-1964	1927-1948
Total weeks	1,266	639
Total cases	4,919	342
Max. cases per year	552	86
Year (max)	1944	1936
Max. cases per week	33	11
Week (max)	1936, wk 03	1936, wk 14
Average cases per year	189	16
95%CI	(128-250)	(6-26)
Average cases per week	4	1
95%CI	(4-4)	(1-1)



**Figure D2,** Number of cases reported for Meningitis per epidemiological week


## Meningococcal Disease



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Meningococcal Disease (Serogroup unspecified and Invasive all serogroups)

Indicator	Texas
Report period	1951-2009
Total weeks	2,602
Total cases	8,896
Max. cases per year	1,325
Year (max)	1953
Max. cases per week	585
Week (max)	1953, wk 02
Average cases per year	
<i>before 1980</i>	177
<i>95%CI</i>	(89-265)
<i>after 1980</i>	130
<i>95%CI</i>	(101-159)
Average cases per week	
<i>before 1980</i>	4
<i>95%CI</i>	(3-5)
<i>after 1980</i>	3
<i>95%CI</i>	(3-3)



**Figure D2**, Number of cases reported for Meningococcal Disease per epidemiological week

## Mumps




**Figure D1**, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1**, Summary information for Mumps

Indicator	Texas	Galveston	San antonio	Houston
Report period	1968-2002	1924-1932	1924-1932	1924-1932
Total weeks	1,684	453	437	404
Total cases	59,479	33	105	190
Max. cases per year	11,236	18	20	85
Year (max)	1968	1925	1928	1927
Max. cases per week	545	5	5	33
Week (max)	1968, wk 14	1928, wk 11	1928, wk 05	1927, wk 05
Average cases per year				
before 1980	4,229	4	12	21
95%CI	(2,161-6,297)	(-1-9)	(7-17)	(1-41)
after 1980	205	-	-	-
95%CI	(132-278)	-	-	-
Average cases per week				
before 1980	83	0	0	0
95%CI	(76-90)	(0-0)	(0-0)	(0-0)
after 1980	4	-	-	-
95%CI	(3-5)	-	-	-

# Preliminary data, not for publication or official use



**Figure D2,** Number of cases reported for Mumps per epidemiological week


## Pneumonia




**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1,** Summary information for Pneumonia (Unspecified)

Indicator	Texas
Report period	1948-1951
Total weeks	169
Total cases	49,355
Max. cases per year	18,432
Year (max)	1950
Max. cases per week	1,026
Week (max)	1950, wk 10
Average cases per year	12,339
95%CI	(1,064-23,614)
Average cases per week	292
95%CI	(260-324)




**Figure D2,** Number of cases reported for Pneumonia per epidemiological week



**Figure D3,** Total number of all cases reported for Pneumonia by each city for the entire time period


## Poliomyelitis



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Poliomyelitis (Total)

Indicator	Texas
Report period	1927-1971
Total weeks	1,968
Total cases	28,727
Max. cases per year	3,903
Year (max)	1952
Max. cases per week	296
Week (max)	1952, wk 28
Average cases per year	
<i>before 1960</i>	831
<i>95%CI</i>	(477-1,185)
<i>after 1960</i>	44
<i>95%CI</i>	(2-86)
Average cases per week	
<i>before 1960</i>	17
<i>95%CI</i>	(16-18)
<i>after 1960</i>	2
<i>95%CI</i>	(2-2)



**Figure D2**, Number of cases reported for Poliomyelitis per epidemiological week


## Rabies in Animals



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1,** Summary information for Rabies in Animals


Indicator	Texas	San antonio
Report period	1948-2009	1952-1953
Total weeks	2,484	72
Total cases	30,212	163
Max. cases per year	1,941	111
Year (max)	2001	1952
Max. cases per week	1,201	7
Week (max)	2001, wk 47	1952, wk 09
Average cases per year	530	82
95%CI	(430-630)	(-293-457)
Average cases per week	12	2
95%CI	(11-13)	(2-2)



**Figure D2,** Number of cases reported for Rabies in Animals per epidemiological week

# Preliminary data, not for publication or official use


## Rubella



**Figure D1**, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1, Summary information for Rubella**

Indicator	Texas
Report period	1966-2002
Total weeks	1,503
Total cases	26,311
Max. cases per year	8,298
Year (max)	1970
Max. cases per week	682
Week (max)	1970, wk 16
Average cases per year	
<i>before 1970</i>	3,223
<i>95%CI</i>	(-853-7,299)
<i>after 1970</i>	329
<i>95%CI</i>	(33-625)
Average cases per week	
<i>before 1970</i>	77
<i>95%CI</i>	(62-92)
<i>after 1970</i>	8
<i>95%CI</i>	(7-9)



**Figure D2**, Number of cases reported for Rubella per epidemiological week


## Salmonellosis



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Salmonellosis (PHLIS and Unspecified)

Indicator	Texas
Report period	1999-2009
Total weeks	487
Total cases	21,319
Max. cases per year	3,609
Year (max)	2003
Max. cases per week	918
Week (max)	2003, wk 42
Average cases per year	1,938
95%CI	(1,438-2,438)
Average cases per week	44
95%CI	(36-52)



**Figure D2,** Number of cases reported for Salmonellosis per epidemiological week


## Scarlet Fever




**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1,** Summary information for Scarlet Fever (Including streptococcal sore throat and Unspecified)

Indicator	Texas	Galveston	San antonio	Houston
Report period	1927-1969	1907-1948	1907-1953	1906-1953
Total weeks	1,835	1,420	1,423	1,498
Total cases	437,657	947	1,838	3,531
Max. cases per year	36,642	64	146	212
Year (max)	1967	1933	1934	1945
Max. cases per week	1,485	9	15	15
Week (max)	1966, wk 08	1934, wk 12	1934, wk 05	1935, wk 05
Average cases per year	11,829	24	50	86
95%CI	(7,057-16,601)	(19-29)	(39-61)	(68-104)
Average cases per week	239	1	1	2
95%CI	(225-253)	(1-1)	(1-1)	(2-2)




**Figure D2,** Number of cases reported for Scarlet Fever per epidemiological week



**Figure D3,** Total number of all cases reported for Scarlet Fever by each city for the entire time period


## Shigellosis



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Shigellosis (Unspecified and PHLIS)

Indicator	Texas
Report period	1999-2009
Total weeks	502
Total cases	23,671
Max. cases per year	5,292
Year (max)	2003
Max. cases per week	1,713
Week (max)	2005, wk 50
Average cases per year	2,152
95%CI	(1,243-3,061)
Average cases per week	47
95%CI	(37-57)



**Figure D2,** Number of cases reported for Shigellosis per epidemiological week


## Streptococcal Disease, Invasive Group A



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Streptococcal Disease, Invasive Group A

Indicator	Texas
Report period	2002-2009
Total weeks	365
Total cases	1,797
Max. cases per year	481
Year (max)	2003
Max. cases per week	72
Week (max)	2003, wk 45
Average cases per year	225
95%CI	(121-329)
Average cases per week	5
95%CI	(4-6)



**Figure D2,** Number of cases reported for Group A Streptococcal Disease per epidemiological week


## Streptococcal Sore Throat



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1, Summary information for Streptococcal Sore Throat**

Indicator	Texas
Report period	1960-1961
Total weeks	102
Total cases	78,950
Max. cases per year	40,666
Year (max)	1960
Max. cases per week	1,533
Week (max)	1960, wk 03
Average cases per year	39,475
95%CI	(24,342-54,608)
Average cases per week	774
95%CI	(722-826)



**Figure D2,** Number of cases reported for Streptococcal Sore Throat per epidemiological week


## Streptococcus Pneumoniae, Invasive Disease



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Streptococcus Pneumoniae, Invasive Disease (Drug resistant <5 years)

Indicator	Texas
Report period	2004-2004
Total weeks	28
Total cases	45
Max. cases per year	45
Year (max)	2004
Max. cases per week	9
Week (max)	2004, wk 34
Average cases per year	45
95%CI	-
Average cases per week	2
95%CI	(1-3)



**Figure D2,** Number of cases reported for Streptococcus Pneumoniae, Invasive Disease per epidemiological week


## Syphilis



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Syphilis (Primary and secondary and Civilian primary and secondary)

Indicator	Texas
Report period	1972-2009
Total weeks	1,794
Total cases	82,616
Max. cases per year	5,875
Year (max)	1982
Max. cases per week	2,120
Week (max)	1991, wk 30
Average cases per year	2,174
95%CI	(1,582-2,766)
Average cases per week	46
95%CI	(43-49)



**Figure D2,** Number of cases reported for Syphilis per epidemiological week


## Tetanus



**Figure D1**, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1, Summary information for Tetanus**

Indicator	Texas
Report period	1962-1981
Total weeks	731
Total cases	308
Max. cases per year	39
Year (max)	1967
Max. cases per week	5
Week (max)	1967, wk 32
Average cases per year	
<i>before 1970</i>	23
<i>95%CI</i>	(16-30)
<i>after 1970</i>	9
<i>95%CI</i>	(7-11)
Average cases per week	
<i>before 1970</i>	1
<i>95%CI</i>	(1-1)
<i>after 1970</i>	0
<i>95%CI</i>	(0-0)



**Figure D2**, Number of cases reported for Tetanus per epidemiological week


## Tuberculosis



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Tuberculosis (Unspecified)

Indicator	Texas
Report period	1975-2005
Total weeks	1,309
Total cases	49,121
Max. cases per year	2,690
Year (max)	1991
Max. cases per week	763
Week (max)	1991, wk 30
Average cases per year	1,585
95%CI	(1,363-1,807)
Average cases per week	38
95%CI	(36-40)



**Figure D2,** Number of cases reported for Tuberculosis per epidemiological week


## Tularemia



**Figure D1**, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1**, Summary information for Tularemia

Indicator	Texas
Report period	1942-1993
Total weeks	1,419
Total cases	988
Max. cases per year	96
Year (max)	1948
Max. cases per week	46
Week (max)	1947, wk 50
Average cases per year	22
95%CI	(15-29)
Average cases per week	1
95%CI	(1-1)



**Figure D2**, Number of cases reported for Tularemia per epidemiological week

## Typhoid Fever




**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Typhoid Fever (Unspecified and Including paratyphoid fever)

Indicator	Texas	Galveston	San antonio	Houston
Report period	1927-2005	1907-1953	1907-1952	1911-1953
Total weeks	3,227	1,593	1,499	1,356
Total cases	23,311	1,172	1,308	1,131
Max. cases per year	1,823	140	172	73
Year (max)	1934	1916	1907	1918
Max. cases per week	414	17	23	16
Week (max)	1953, wk 02	1930, wk 46	1923, wk 26	1930, wk 47
Average cases per year				
before 1950	828	30	37	36
95%CI	(616-1,040)	(18-42)	(22-52)	(30-42)
after 1950	71	1	1	8
95%CI	(29-113)	-	-	(-17-33)
Average cases per week				
before 1950	17	1	1	1
95%CI	(16-18)	(1-1)	(1-1)	(1-1)
after 1950	2	1	1	1
95%CI	(1-3)	-	-	(1-1)

Preliminary data, not for publication or official use




**Figure D2,** Number of cases reported for Typhoid Fever per epidemiological week



**Figure D3,** Total number of all cases reported for Typhoid Fever by each city for the entire time period


## Typhus Fever



**Figure D1**, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1**, Summary information for Typhus Fever

Indicator	Texas
Report period	1942-1961
Total weeks	573
Total cases	8,328
Max. cases per year	1,761
Year (max)	1945
Max. cases per week	84
Week (max)	1945, wk 46
Average cases per year	490
95%CI	(157-823)
Average cases per week	15
95%CI	(14-16)



**Figure D2**, Number of cases reported for Typhus Fever per epidemiological week


## West Nile Disease



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for West Nile Disease (Neuroinvasive, Non-neuroinvasive)

Indicator	Texas
Report period	2003-2009
Total weeks	176
Total cases	1,041
Max. cases per year	438
Year (max)	2003
Max. cases per week	127
Week (max)	2003, wk 42
Average cases per year	149
95%CI	(19-279)
Average cases per week	6
95%CI	(4-8)



**Figure D2,** Number of cases reported for West Nile Disease per epidemiological week

## Whooping Cough




**Figure D1**, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1**, Summary information for Whooping Cough

Indicator	Texas	Galveston	San antonio	Houston
Report period	1937-2009	1924-1948	1907-1953	1924-1953
Total weeks	2,251	1,139	1,225	1,158
Total cases	189,921	256	1,213	1,277
Max. cases per year	21,365	88	226	219
Year (max)	1947	1942	1940	1943
Max. cases per week	1,482	18	23	29
Week (max)	2005, wk 50	1924, wk 24	1940, wk 32	1937, wk 26
Average cases per year				
<i>before 1960</i>	9,384	10	40	47
95%CI	(7,231-11,537)	(1-19)	(20-60)	(23-71)
<i>after 1960</i>	323	-	-	-
95%CI	(144-502)	-	-	-
Average cases per week				
<i>before 1960</i>	194	0	1	1
95%CI	(185-203)	(0-0)	(1-1)	(1-1)
<i>after 1960</i>	9	-	-	-
95%CI	(7-11)	-	-	-

Preliminary data, not for publication or official use



**Figure D2,** Number of cases reported for Whooping Cough per epidemiological week

## Diseases with fragmented data over time

For a number of diseases, only fragmented data was available for Texas. This was due to the inclusion of these diseases in the weekly surveillance system for short periods or widely dispersed periods of time. Due to these wide gaps or limited data, data for these diseases were not presented in separate sections but in the table below. Work will continue to include more data in the Tycho database and complete data sets for each disease where possible. Collaborations with local, state and federal public health agencies will be made to explore data availability and opportunities to include these in the Tycho database.

**Table 2**, Summary information on the occurrence of diseases with fragmented data in Texas

Disease	Report type	Report period	Number of reports	Total cases
Anthrax	State	1942-1945	142	9
Botulism	State	1952-1952	1	1
Brucellosis [undulant fever]	City	1952-1952	1	1
Brucellosis [undulant fever]	State	1943-1982	816	10388
Dengue	City	1924-1925	14	0
Dysentery	State	1942-1948	333	37762
Encephalitis	City	1941-1953	1351	44
Encephalitis	State	1942-1994	1607	5193
Giardiasis	State	2003-2004	13	4
Leprosy	City	1924-1924	1	0
Leprosy	State	1942-1991	509	197
Pellagra	City	1924-1932	835	287
Psittacosis	State	1956-1961	22	67
Rocky mountain spotted fever	City	1953-1953	2	4
Rocky mountain spotted fever	State	1942-2009	1407	857
Toxic shock syndrome	State	1984-1992	186	87
Yellow fever	City	1907-1907	1	1

Preliminary data, not for publication or official use

## **Project Tycho**

This report provides preliminary data for the state of Texas available in the Tycho database. This database is currently being beta tested and these data cannot be used for publication or other official use at this time. An open access release to the general public is planned for later in 2011.

Please visit the Tycho website for more information and to query the database at: [www.tycho.pitt.edu](http://www.tycho.pitt.edu). For further information regarding the Tycho project, contact Dr. Wilbert van Panhuis at the University of Pittsburgh Graduate School of Public Health.

Dr. Wilbert van Panhuis, MD PdD  
Graduate School of Public Health  
University of Pittsburgh  
130 DeSoto Street  
704 Parran Hall  
Pittsburgh, 15261 PA  
Tel: 412-624-7693  
Email: wav10@pitt.edu