

# Project Tycho

*Preliminary data for the state of Illinois*

## Tycho database beta test version

*The data presented in this report are of preliminary nature and should not be used for publication or other types of official use*

June 2011

University of Pittsburgh Graduate School of Public Health



University of Pittsburgh



## Table of Content

<b>Acknowledgements</b>	<b>page 3</b>
<b>Contact information</b>	<b>page 4</b>
<b>Introduction</b>	<b>page 5</b>
<b>Vision and goal</b>	<b>page 7</b>
<b>Collaboration</b>	<b>page 8</b>
<b>Summary of methods</b>	<b>page 9</b>
<b>Overview of data available for the state of Illinois</b>	<b>page 12</b>
<b>Disease specific data for Illinois</b>	<b>page 19</b>
- AIDS	page 19
- Chickenpox	page 20
- Chlamydia	page 22
- Cryptosporidiosis	page 23
- Diphtheria	page 24
- Escherichia Coli	page 27
- Giardiasis	page 28
- Gonorrhea	page 29
- Haemophilus Influenzae	page 30
- Hepatitis	page 31
- Influenza	page 33
- Legionellosis	page 36
- Malaria	page 37
- Measles	page 38
- Meningitis	page 41
- Meningococcal disease	page 43
- Mumps	page 45
- Pellagra	page 47
- Pneumonia	page 48
- Poliomyelitis	page 51
- Rabies in animals	page 54
- Rubella	page 55
- Salmonellosis	page 56
- Scarlet fever	page 57
- Shigellosis	page 60
- Streptococcal disease, invasive group A	page 61
- Streptococcal sore throat	page 62
- Streptococcus pneumonia, invasive disease	page 63

## Disease specific data for Illinois, continued

- Syphilis	page 64
- Tuberculosis	page 65
- Typhoid fever	page 68
- Whooping cough	page 71

Diseases with fragmented data	page 73
-------------------------------	---------

## Acknowledgements

The development of this project would not have been possible without data collected every week by staff from city-, county-, state- and federal public health departments. We would like to acknowledge their invaluable contribution to the overall body of surveillance data that were digitized during this project. Particularly, we acknowledge staff from the Council of State and Territorial Epidemiologists (CSTE) and the US Centers for Disease Control for their overall efforts in disease surveillance and their particular input in this project.

We also acknowledge the help of many contributors that have contributed to the digitization and processing of weekly surveillance data included in this database. Digital Divide Data: Linda Thomas, Eric Gold, Heak Hok, Villa Kem and Hetel Patel. University of Pittsburgh Graduate School of Public Health: Steve Wisniewsky, Heather Eng, Sharon Crow, David Galloway, Jia Xu and Raaka Kumbhakar. University of Pittsburgh School of Information Sciences: Vladimir Zadorozhny, Ying-Feng Hsu, David Paul Koenig, Fatimah Ridwan and Divyasheel Sharma. Johns Hopkins Bloomberg School of Public Health: Derek Cummings. Pittsburgh Supercomputing Center: David Graham and Andrew Adams.

This project was made possible with funding from the Bill & Melinda Gates Foundation through the Vaccine Modeling Initiative (grant ID 49276).

The Tycho development team at the University of Pittsburgh Graduate School of Public Health:

Willem G. van Panhuis

John Grefenstette

Shawn T. Brown

Su Yon Jung

Nian Shong Chok

Anne Cross

Donald S. Burke

## Contact information

The Tycho database is currently being tested and a login account will be required to use the database and website ([www.tycho.pitt.edu](http://www.tycho.pitt.edu)). All State Epidemiologists, CDC partners, other federal agencies and our research collaborators will be provided with user accounts. User accounts can also be provided upon specific request. Note that the quality of the data during the testing phase cannot be guaranteed to be sufficient for publication or official use.

**Point of contact for the Tycho project:**

Dr. Willem G. van Panhuis, MD PhD  
Graduate School of Public Health  
University of Pittsburgh  
130 DeSoto Street  
704 Parran Hall  
Pittsburgh PA, 15261  
Email: [wav10@pitt.edu](mailto:wav10@pitt.edu)  
Tel: 412-624-7693

## Introduction

**Project Tycho** started in 2009 as part of the Vaccine Modeling Initiative (VMI) at the University of Pittsburgh Graduate School of Public Health, funded by the Bill & Melinda Gates foundation. This project aims to contribute to the availability of public health data for research and policy making. The vision for this project as described in the next section is a central access point for public health data of a detailed spatial and temporal scale for all countries. Currently, the Tycho database contains 10 million records that each represent a weekly report from a location for a specific disease. These records were extracted from weekly US Nationally Notifiable Disease Surveillance Reports between 1888 and 2009 (6300 reports) using double data entry (200 million keystrokes). The current database includes a total of 90 million reported cases and 4 million reported deaths due to notifiable diseases in the United States for the last 122 years.

Project **Tycho** is named after the Danish nobleman Tycho Brahe (1546 – 1601), who meticulously collected astronomical data. After Brahe's death, his assistant Kepler used these data to develop area laws of planetary motion.

## Background

***"No health department, State or local, can effectively prevent or control diseases without knowledge of when, where and under what conditions cases are occurring"***

(Quote stated above weekly surveillance reports published in the Public Health Reports, 1888-1951)

This quote has accompanied weekly reports on the prevalence of notifiable diseases in the United States as long as these were published in the Public Health Reports, emphasizing the importance of data for disease control policy making. In the current day and age, this statement has lost none of its significance. On the contrary, as infectious diseases continue to pose challenges to the global public health system in an increasingly interconnected global society. The public health system has responded to this challenge by evermore sophisticated disease surveillance systems. The availability of internet and information technology has facilitated collection of detailed data on infectious disease incidence, the spread of pathogens, disease determinants, health behavior, etc. New technologies such as remote sensing and mobile reporting systems have advanced measurement of these factors in both high and low resource settings. The enormous wealth of public health information that is currently being accrued also poses its own challenges for data storage, management, preservation and dissemination. Public health agencies will need to allocate resources and develop capacity to specifically address these issues.

Although the importance of public health data collection has been widely recognized, the dissemination of data has been neglected for many reasons. The above quote does not only imply data collection but also its use and dissemination as knowledge on disease occurrence can only be derived from data after processing and analysis. Not only has disease surveillance become more sophisticated, analytical tools have changed as well and now include computational models to assess disease transmission and to evaluate disease control options. These methods have introduced the opportunity for data intensive studies of fine spatial-temporal patterns using dynamic models. Disease surveillance data at such scale are often not available for analysis for multiple reasons. First, surveillance data are not collected for research purposes but for disease monitoring and planning. Secondly, public health staff often lack time or capacity to prepare these data for analysis. Third, surveillance methodology often lacks transparency and standardization, reducing options for scientific analyses. Fourth, disease surveillance data are often

kept confidential and ownership or data use requirements are often unclear. Finally, archiving and preservation of surveillance data are often not a priority and disaggregated records may get lost, losing the opportunity to ever use these data for analysis.

The Tycho project aims at overcoming some of these challenges by providing open access to large public health datasets of great spatial-temporal detail. Open access to these data will provide great opportunities for scientific analysis and better informed policy making on disease control.

## **The Tycho database and website**

As described in the methods section, a large heterogeneity in reporting practice was observed over the past 122 years of weekly disease surveillance and substantial efforts were made to clean and standardize these data. The current online database includes all cleaned and standardized data in a format that allows comparisons across time and space. It does not include parts of the data that require additional cleaning and research such as classification of disease subcategories.

Each record in the database provides a reported number of cases or deaths due to a specific disease for a specific location, for a certain week. The database can be searched online ([www.tycho.pitt.edu](http://www.tycho.pitt.edu)) by selecting a dataset, an aggregation method and an output format. Data can be viewed in tables, graphics and maps and these can be downloaded for further use.

## **Data availability**

The data provided by the Tycho database reflect the weekly US Nationally Notifiable Disease Surveillance System between 1888 and 2009. It includes all diseases that were reported as part of this system, but it does not yet include records from other surveillance systems such as the annual system or disease specific subsystem (such as the influenza surveillance system).

The available data also reflects changes in reporting practices over time. This explains why city or state level reports are only available for a certain time period (1888-1953 for cities and 1927-2009 for states) and why morbidity and mortality reports are available for different time periods. We are currently working on inclusion of additional data as described in the section on collaboration.

## **Testing and release of the Tycho database**

The Tycho database is currently in a beta testing phase that will start during the 2011 annual meeting of the Council of State and Territorial Epidemiologists (CSTE) in Pittsburgh June 12-16. This phase will end with a release to the general public in the fall of 2011. During the testing phase, invited users will be provided with login accounts that will enable full use of all features of the website and database. Invited users will include all state epidemiologists, partners in the CDC and other federal agencies as well as research collaborators. During this phase, data cleaning and standardization will continue. The quality of the data will not be sufficient for publication or official use during this phase.

## **Future developments**

After optimization of the database and website during the testing phase, the first version of the Tycho database will be released to the general public. After that, efforts to clean and standardize the data will continue and more data will be released in subsequent versions. In addition, ongoing and new collaborations for inclusion of new data will allow gradual expansion of the dataset over time and increased opportunities for data driven analysis and policy making.

## Vision and goal

The vision for project Tycho is based on the value of open access to detailed, disaggregated public health data for scientific analysis and policy making. Starting with 122 years of weekly US surveillance data, it is our hope that this example will convince public health authorities worldwide of the value of this resource and will stimulate similar contributions from other countries.

In the past, data sharing and archiving have not been a priority in public health or among health sciences in general. Continuous and diligent public health data collection has led to a large pool of disaggregated data scattered over the world including the internet, ranging from paper archives to basements of local health departments. The need for open access to a central repository of public health data has been recognized and both benefits and challenges of open access to data in general have been well described. Open access to data could lead to increased accountability, transparency, innovation, collaboration, cost-effectiveness, replication of results, the development of new methods and insights, and a narrower data access gap between low- and higher income countries. Ultimately, this would result in the advancement of science and technology to the benefit of all.

Challenges to open access include inconsistent formats, lack of annotation or metadata, lack of information technology (IT) capacity, a lack of incentive, and cultural norms around data ownership. Open access to public health data in particular is confronted by decentralized archiving of disease reports, lack of time and capacity for data management, degraded physical format of records and archives, and governmental level legal constraints. Multiple principles and requirements for a central open access repository for public health data have been described and include (1) capacity building for data management in low- and middle income countries, (2) feedback loops to data contributors, (3) sustainability, (4) common standards, (5) interoperability, and (6) user-friendliness.

The ultimate goal of the Tycho project is to provide a central global public health data access point. Historical as well as current public health data are of great value if archived and accessible for research and analysis. Open access will enable the use of analytical capacity from around the globe which will lead to new discoveries of disease patterns and control policies. Furthermore, a central data access point will facilitate data archiving and preservation into the future which will be an increasing need in a data rich public health environment.

## Collaboration

The application of the surveillance data from the Tycho database as well as the inclusion of new data will require a wide range of collaborations with public health offices in the US and abroad. The Tycho project has currently made the first step of digitizing 122 years of weekly US surveillance reports that had already been published. The next step will be the completion of the existing dataset as well as the inclusion of new data from unpublished sources.

This report provides an overview of data available from the Tycho database for the state of Illinois. This includes all data that has ever been published at state or city level for Illinois in the weekly US Nationally Notifiable Disease Surveillance System. As you will find, the data availability will vary greatly over time, between city and state level reports, between diseases and between morbidity vs. mortality reports.

The Tycho development team would look forward to work with state health departments in the US to:

1. Better understand the current Tycho data for each state (eg. by collecting historic documentation)
2. Provide better usability and applications of current Tycho data for each state by continued joint development of website and database features.
3. Collect and/or digitize official, confirmed data to validate the current (preliminary) weekly data.
4. For each state, collect and/or digitize new data that has not yet been included. For example all city level reports discontinued in 1953 and completion of city level data until 2009 would greatly increase opportunities for analysis of disease patterns and trends.
5. Provide support to state health offices to manage, preserve and provide access to public health data.



## Summary of methods

Detailed documentation on the methods used for the Tycho database have been provided on the website ([www.tycho.pitt.edu](http://www.tycho.pitt.edu)). This section describes these methods in short.

### Data collection

Weekly reports that contain tables on the occurrence of nationally notifiable diseases have been published since 1888 by public health authorities at the Federal level in various journals. Table 1 provides the list of publications and the responsible agency since 1888. All weekly nationally notifiable disease reports between 1888 and 1951 could be retrieved from the PubMed Central repository of the National Library of Medicine <sup>1</sup>. For 1995 to 2009, these reports could be retrieved from the MMWR digital archive on the CDC website <sup>2</sup>. Most weekly reports between 1952 and 1995 could be retrieved from the HathiTrust Digital Library <sup>3 4</sup>, but many could not be found and had to be copied from hard copies of MMWR issues in the University of Pittsburgh library.

Table 1, publications and responsible Federal agencies for nationally notifiable disease reports

Time period	Publication title	Responsible federal agency
1888-1889	Weekly Abstract of Sanitary Reports	US Marine Hospital Service
1890-1895	Abstract of Sanitary Reports	US Marine Hospital Service
1896-1901	Public Health Reports	US Marine Hospital Service
1902-1911	Public Health Reports	US Public Health and Marine Hospital Service
1912-1951	Public Health Reports	US Public Health Service
1952-1960	Morbidity and Mortality Weekly Report	National Office of Vital Statistics, US Public Health Service
1961-1969	Morbidity and Mortality Weekly Report	Communicable Disease Center
1970-1991	Morbidity and Mortality Weekly Report	Center for Disease Control
1992-2009	Morbidity and Mortality Weekly Report	Centers for Disease Control and Prevention

### Inclusion criteria

Weekly reports of each year were reviewed systematically to assess the diseases reported. We included all tables that provided disease specific information by week for US cities, townships, counties or states. Tables that provided summary or aggregated information by month, year or at the national level were not included. Similarly, tables that did not contain disease specific information (such as all cause mortality) were not included.

### Data entry

Weekly reports were downloaded or scanned as PDF files and selected tables with notifiable disease reports were entered into computer spreadsheets in a highly standardized fashion using double data entry. During the second round of data entry, operators could not see what had been entered in the first round and could not continue if the system detected a discrepancy between the second and first entry

<sup>1</sup> <http://www.pubmedcentral.nih.gov/tocrender.fcgi?journal=333&action=archive>

<sup>2</sup> <http://www.cdc.gov/mmwr>

<sup>3</sup> <http://catalog.hathitrust.org/Record/003910026>

<sup>4</sup> <http://catalog.hathitrust.org/Record/003843660>

for a specific value. Such discrepancies could only be resolved by checking the PDF file and try again or discussion with the group leader and verification of a value in the source documents.

## **Quality control for data entry**

The accuracy of data entry was checked at various levels. First, completeness of data was verified by comparing the content of entered data with PDF sources files. Secondly, accuracy of data entry was verified by multiple rounds of comparing random samples of entered files with PDF source files. Thirdly, data formatting was verified by various checks to ensure appropriate formatting for data loading.

## **Data loading and standardization**

All data was entered in Excel spreadsheets and various components of these spreadsheets were loaded in data files. Table titles, column headers, place names and reported numbers were loaded in separate files. These files were used to extract information on each reported number, including:

1. the disease reported
2. the disease subcategory reported
3. cases or deaths reported
4. the reporting location (name, state and type of location)
5. the time period for which a number was reported
6. the date of publication of the original weekly report associated with a reported number

## **Integration**

All reported numbers and extracted information was integrated in one database with one record per reported number and associated information.

## **Post-processing quality control**


After integration of all data in one database, checks were performed to detect duplicate reports and data inconsistencies. Duplicate records were removed and inconsistencies resolved by verification with original PDF source files.

## **Data filtering**

The digitized version of all historical weekly US nationally notifiable disease surveillance records is a very heterogeneous dataset. It took a substantial number of data processing protocols to standardize time and space variables as well as disease names. All standardized records have been separated from non-standard records by a filter. All remaining heterogeneity in non-standard records is inherent to the surveillance system and can only be standardized after further analysis will have been completed (eg. remaining heterogeneity in reporting periods for reports before 1953 and standardization of disease subcategories). The largest proportion of data (>4 million records) has been standardized however and has been made available in the current testing version.

## **Data visualization**

Figures in this reports were made with the R system, version 2.9.2 and the maps were generated by the GAIA platform developed at the University of Pittsburgh Graduate School of Public Health in collaboration with the Pittsburgh Supercomputing Center (PSC). See <http://midas-pitt.psc.edu/gaia> for more information.



**Figure 1, Schematic of the Tycho data architecture.** Access to a consistent, standardized subset of the digitized US weekly surveillance reports will be provided for beta testing through a website that will allow querying and downloading of data. Data from the development database will be pushed through a filter to feed the Tycho database.

The current Tycho database will provide the reported number of cases or deaths reported by city or state health authorities to the federal health agency for all weeks between 1888 and 2009. Currently, only standardized, consistent data are being released for testing, according to the following criteria:

1. Only weekly reports are provided (reports for other periods such as 10 days, 2 weeks, 1 month, 1 year) are currently not included due to lack of comparability with the weekly reports.
2. Only reports that were published in the same year as the reporting period were included (this excludes updates or comparisons that were published more than a year after the original reporting period)
3. Only numbers were included for which information about the location, time period and disease could be extracted.
4. Only numbers or time periods for which no disease subcategories were reported were included.

Each of the steps described above have been described in detailed protocols that are posted in the documentation part of the Tycho website ([www.tycho.pitt.edu](http://www.tycho.pitt.edu)).

## Website

The Tycho website features simple and advanced searches of the data, visualizations and downloading of machine readable files. A dataset can be selected by specifying: 1) the disease of interest, 2) the location(s) of interest and 3) the time period. In a second step, the type of aggregation and length of increments can be specified and in a third step, the type of output (table, graphic or map) can be selected. After output has been generated, data can be downloaded.

As mentioned above, data availability varies largely by disease, time period and location. We aimed to restrict selection options by data availability to avoid "no results available".

Access to the most of the website will be password protected during the testing phase. For invited users after login, full functionality of the database and website will be available. During this phase however, the data quality will not be sufficient for publication or official use.

## Overview of data available for the state of Illinois

In this section, an overview of data available for the state of Illinois will be provided. As described earlier, data availability depends entirely on historical reporting practices in the weekly US Nationally Notifiable Disease Surveillance System between 1888 and 2009. For this report, we only included morbidity reports (cases) to improve the format and limit the size of this report. As mentioned earlier, the analysis and standardization of has not been completed yet for all records in the database and preliminary data are provided here that may not yet be available in the online database.

This section will provide a general overview of data availability. The next sections will provide disease specific data from state and city level reports of Illinois. Data for a maximum of 3 major cities were provided depending on availability (city data was only reported until 1953). Some summary data will be provided at the end of this report, for a subset of diseases for which data was fragmented over time.

Table 1 lists the number of weekly state or city reports that are available for Illinois per each disease and subcategory.

**Table 1, Number of weekly state or city reports per disease and subcategory**

<b>Disease</b>	<b>City</b>	<b>State</b>
Aids	-	934
Anthrax	-	143
Botulism	1	-
Brucellosis [undulant fever]	27	756
Chickenpox [varicella]	450	565
Chlamydia	-	654
Cryptosporidiosis	-	437
Dengue	8	-
Diphtheria	2157	1708
Dysentery		
<i>Amebic</i>	-	324
<i>Bacillary</i>	-	270
<i>Unspecified</i>	-	196
Encephalitis		
<i>Lethargic</i>	441	-
<i>Post infectious</i>	-	621
<i>Primary [infectious] including unspecified</i>	362	1608
Escherichia coli		
<i>EHEC 0157</i>	-	176
<i>EHEC non-0157</i>	-	62
<i>EHEC non serogrouped</i>	-	34
<i>O157:H7 NETSS</i>	-	270
<i>O157:H7 PHLIS</i>	-	160
<i>STEC</i>	-	153

**Table 1,** Number of weekly state or city reports per disease and subcategory, *continued*

<b>Disease</b>	<b>City</b>	<b>State</b>
Giardiasis	-	367
Gonorrhea		
<i>Civilian</i>	-	754
<i>Unspecified</i>	-	1056
Haemophilus influenzae		
<i>Age &lt;5 non-serotype B</i>	-	36
<i>Age &lt;5 unknown serotype</i>	-	89
<i>All ages all serotypes</i>	-	589
Hepatitis		
<i>Acute type A</i>	-	371
<i>Acute type B</i>	-	315
<i>Acute type C</i>	-	85
<i>Acute type NA NB [including C]</i>	-	41
<i>All types, &lt;20 years</i>	-	204
<i>All types, &gt;=20 years</i>	-	104
<i>All types, all ages</i>	51	708
<i>Type A [infectious]</i>	1	1641
<i>Type B [serum]</i>	1	1457
<i>Type NA NB [including C]</i>	-	677
<i>Type unspecified</i>	1	719
Influenza	1263	1166
Legionellosis	-	739
Leprosy	1	387
Listeriosis	-	133
Lyme disease	-	424
Malaria		
<i>Civilian</i>	-	3
<i>Military</i>	-	10
<i>Unspecified</i>	2	1771
Measles		
<i>Imported</i>	-	480
<i>Indigenous</i>	-	597
<i>Unspecified</i>	2231	2964
Meningitis		
<i>Aseptic</i>	1	1221
<i>Meningococcus</i>	1023	1280
<i>Unspecified</i>	657	5

**Table 1,** Number of weekly state or city reports per disease and subcategory, *continued*


<b>Disease</b>	<b>City</b>	<b>State</b>
Meningococcal disease		
<i>All serogroups</i>	-	29
<i>Invasive all serogroups</i>	-	178
<i>Invasive serogroup unknown</i>	-	45
<i>Serogroup unspecified</i>	82	2430
Mumps	450	1651
Pellagra	438	-
Pneumonia		
<i>Lobar</i>	21	-
<i>Unspecified</i>	605	170
Poliomyelitis		
<i>Non paralytic</i>	-	201
<i>Paralytic</i>	-	482
<i>Total</i>	1572	1861
Psittacosis	-	48
Rabies in animals	1	2451
Rocky mountain spotted fever	3	1219
Rubella	-	1515
Salmonellosis		
<i>NETSS</i>	-	125
<i>PHLIS</i>	-	116
<i>Unspecified</i>	-	373
Scarlet fever		
<i>Including streptococcal sore throat</i>	97	564
<i>Unspecified</i>	2155	1284
Shigellosis		
<i>NETSS</i>	-	125
<i>PHLIS</i>	-	116
<i>Unspecified</i>	-	373
Streptococcal disease, invasive group a	-	373
Streptococcal sore throat	-	102
Streptococcus pneumoniae invasive disease		
<i>Drug resistant &lt;5 years</i>	-	77
<i>Drug resistant A</i>	-	5
<i>Drug resistant all ages</i>	-	187
<i>Drug resistant B</i>	-	5
<i>Non drug resistant &lt;5 years</i>	-	179
Syphilis		
<i>Civilian primary and secondary</i>	-	728
<i>Congenital</i>	-	147
<i>Primary and secondary</i>	-	1088

**Table 1,** Number of weekly state or city reports per disease and subcategory, *continued*

<b>Disease</b>	<b>City</b>	<b>State</b>
Tetanus	-	489
Toxic shock syndrome	-	265
Trichiniasis	4	8
Tuberculosis [phthisis pulmonalis]		
<i>New active</i>	-	144
<i>Unspecified</i>	875	1485
Tularemia	4	1060
Typhoid fever [enteric fever]		
<i>Including paratyphoid fever</i>	338	668
<i>Unspecified</i>	1775	2316
Typhus fever		
<i>Endemic</i>	-	22
<i>Unspecified</i>	41	178
West nile disease		
<i>Neuroinvasive</i>	-	120
<i>Non-neuroinvasive</i>	-	77
Whooping cough [pertussis]	1548	2268

The diseases included in the weekly US Nationally Notifiable Disease Surveillance System varied largely over time and reflected the historical social-political priorities of each time period. Note that the diseases in the weekly system were a subset of all diseases included in the annual Notifiable Disease Surveillance System (that was not entered as part of this project).

The number of locations reporting diseases also varied over time depending on the type of reports submitted. Figure 1 on the next page provides an overview of the number of unique locations reporting each disease per week between 1888 and 2009. For the remainder of this report, the state of Illinois was considered a unique reporting location in addition to the individual city reports. Figure 1 indicates that many locations (mostly cities) were included in the reporting system before 1925, after which the number of cities dropped substantially. After 1953, only state reports were included and city reports were discontinued. These patterns were observed nationwide.




**Figure 1,** The number of locations (in color code) reporting on each disease per week between 1888 and 2009

For the majority of the time period between 1888 and 2009, only state level reports were available. It would be of great interest to complete the collection of city data after 1953 and we will seek collaboration with state health offices for this.


The map in figure 2 (next page) illustrates the geographical distribution of the available city data. It displays the number of weekly reports for any disease available per location that was included (and for which coordinates could be derived)





**Figure 2,** This map displays the number of weekly reports available for any disease per city (note that city reports were only available until 1953)


Figure 3 below provides an overview of the number of diseases for which a weekly report was included at the state level and for individual cities. All cities for which at least 100 weekly reports are available have been listed. For most states, it shows a major drop in the number of cities after 1925 and after 1953, city reports were discontinued except for New York City, which became its own reporting jurisdiction.



**Figure 3,** The number of diseases (in color code) that were reported for the state and city level (state on top row) for each week between 1888 and 2009

## Disease specific data for Illinois


### AIDS



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for AIDS

Indicator	Illinois
Report period	1984-2005
Total weeks	934
Total cases	28,716
Max. cases per year	2,848
Year (max)	1994
Max. cases per week	752
Week (max)	1993, wk 15
Average cases per year	1,305
95%CI	(988-1,622)
Average cases per week	31
95%CI	(26-36)



**Figure D2,** Number of cases reported for AIDS per epidemiological week


## Chickenpox



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Chickenpox

Indicator	Chicago	Springfield
Report period	1924-1932	1924-1932
Total weeks	450	418
Total cases	39,750	2,018
Max. cases per year	5,500	330
Year (max)	1924	1926
Max. cases per week	261	38
Week (max)	1928, wk 48	1924, wk 15
Average cases per year		
before 1995	4,417	224
95%CI	(3,862-4,972)	(182-266)
after 1995	-	-
95%CI	-	-
Average cases per week		
before 1995	88	5
95%CI	(83-93)	(4-6)
after 1995	-	-
95%CI	-	-



**Figure D2,** Number of cases reported for Chickenpox per epidemiological week


## Chlamydia



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Chlamydia

Indicator	Illinois
Report period	1996-2009
Total weeks	653
Total cases	421,910
Max. cases per year	50,269
Year (max)	2007
Max. cases per week	7,060
Week (max)	1999, wk 44
Average cases per year	30,136
95%CI	(22,276-37,996)
Average cases per week	646
95%CI	(594-698)



**Figure D2,** Number of cases reported for Chlamydia per epidemiological week


## Cryptosporidiosis



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Cryptosporidiosis

Indicator	Illinois
Report period	1999-2009
Total weeks	437
Total cases	1,583
Max. cases per year	436
Year (max)	2001
Max. cases per week	372
Week (max)	2001, wk 44
Average cases per year	144
95%CI	(70-218)
Average cases per week	4
95%CI	(2-6)



**Figure D2,** Number of cases reported for Cryptosporidiosis per epidemiological week

## Diphtheria



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1,** Summary information for Diphtheria

Indicator	Illinois	Chicago	Springfield
Report period	1927-1975	1906-1953	1907-1948
Total weeks	1,708	2,129	1,676
Total cases	53,350	155,937	2,409
Max. cases per year	8,512	10,297	341
Year (max)	1929	1917	1915
Max. cases per week	616	526	34
Week (max)	1970, wk 44	1909, wk 09	1915, wk 42
Average cases per year			
before 1940	3,454	4,408	72
95%CI	(1,934-4,974)	(3,419-5,397)	(47-97)
after 1940	151	165	2
95%CI	(52-250)	(4-326)	(0-4)
Average cases per week			
before 1940	71	89	2
95%CI	(67-75)	(86-92)	(2-2)
after 1940	5	4	0
95%CI	(4-6)	(3-5)	(0-0)






Figure D2, Number of cases reported for Diphtheria per epidemiological week





Figure D3, The total number of cases reported for Diphtheria by each city for the entire time period


## Escherichia Coli



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Escherichia Coli (O157:H7 PHLIS, EHEC O157, and STEC)

Indicator	Illinois
Report period	1996-2009
Total weeks	489
Total cases	1,184
Max. cases per year	159
Year (max)	2002
Max. cases per week	68
Week (max)	1996, wk 36
Average cases per year	85
95%CI	(57-113)
Average cases per week	2
95%CI	(1-3)



**Figure D2,** Number of cases reported for Escherichia Coli per epidemiological week


## Giardiasis



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Giardiasis

Indicator	Illinois
Report period	2002-2009
Total weeks	366
Total cases	4,553
Max. cases per year	694
Year (max)	2003
Max. cases per week	228
Week (max)	2004, wk 34
Average cases per year	569
95%CI	(474-664)
Average cases per week	12
95%CI	(10-14)



**Figure D2,** Number of cases reported for Giardiasis per epidemiological week


## Gonorrhea



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Gonorrhea (Unspecified and Civilian)

Indicator	Illinois
Report period	1972-2009
Total weeks	1,809
Total cases	993,754
Max. cases per year	57,880
Year (max)	1975
Max. cases per week	9,427
Week (max)	1999, wk 44
Average cases per year	26,151
95%CI	(21,836-30,466)
Average cases per week	549
95%CI	(521-577)



**Figure D2,** Number of cases reported for Gonorrhea per epidemiological week


## Haemophilus Influenzae



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Haemophilus Influenzae (All ages all serotypes)

Indicator	Illinois
Report period	1996-2009
Total weeks	589
Total cases	890
Max. cases per year	142
Year (max)	2009
Max. cases per week	22
Week (max)	2001, wk 48
Average cases per year	64
95%CI	(45-83)
Average cases per week	2
95%CI	(2-2)



**Figure D2,** Number of cases reported for Haemophilus Influenzae per epidemiological week


## Hepatitis



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1,** Summary information for Hepatitis (Type A [Infectious], Acute type A, Type B [Serum], and Acute type B)


Indicator	Illinois
Report period	1966-2009
Total weeks	2,036
Total cases	26,576
Max. cases per year	1,929
Year (max)	1970
Max. cases per week	934
Week (max)	1970, wk 44
Average cases per year	
before 1990	830
95%CI	(649-1,011)
after 1990	307
95%CI	(212-402)
Average cases per week	
before 1990	18
95%CI	(16-20)
after 1990	7
95%CI	(6-8)



**Figure D2,** Number of cases reported for Hepatitis per epidemiological week




## Influenza




**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1,** Summary information for Influenza

Indicator	Illinois	Chicago
Report period	1919-1951	1920-1948
Total weeks	1,166	1,248
Total cases	59,839	44,712
Max. cases per year	10,721	19,776
Year (max)	1928	1920
Max. cases per week	3,251	9,955
Week (max)	1920, wk 02	1920, wk 04
Average cases per year	2,216	1,542
95%CI	(1,208-3,224)	(157-2,927)
Average cases per week	51	36
95%CI	(40-62)	(18-54)




**Figure D2,** Number of cases reported for Influenza per epidemiological week



**Figure D3, Total number of cases reported for Influenza by each city for the entire time period**


## Legionellosis



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Legionellosis

Indicator	Illinois
Report period	1982-2009
Total weeks	739
Total cases	575
Max. cases per year	98
Year (max)	2009
Max. cases per week	35
Week (max)	2009, wk 39
Average cases per year	23
95%CI	(12-34)
Average cases per week	1
95%CI	(1-1)



**Figure D2,** Number of cases reported for Legionellosis per epidemiological week


## Malaria



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Malaria (Unspecified, Military, and Civilian)

Indicator	Illinois
Report period	1952-2009
Total weeks	1,771
Total cases	1,661
Max. cases per year	185
Year (max)	1969
Max. cases per week	35
Week (max)	1996, wk 48
Average cases per year	33
95%CI	(24-42)
Average cases per week	1
95%CI	(1-1)



**Figure D2,** Number of cases reported for Malaria per epidemiological week

## Measles



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1,** Summary information for Measles (Unspecified)

Indicator	Illinois	Chicago	Springfield
Report period	1927-2001	1906-1953	1907-1948
Total weeks	2,964	2,220	1,399
Total cases	859,887	440,838	18,678
Max. cases per year	88,011	37,769	2,075
Year (max)	1938	1938	1942
Max. cases per week	6,933	3,478	311
Week (max)	1938, wk 09	1938, wk 10	1942, wk 12
Average cases per year before 1970	19,080	9,796	467
95%CI	(13,723-24,437)	(7,534-12,058)	(263-671)
after 1970	1,018	-	-
95%CI	(401-1,635)	-	-
Average cases per week before 1970	383	199	13
95%CI	(355-411)	(185-213)	(11-15)
after 1970	26	-	-
95%CI	(23-29)	-	-






Figure D2, Number of cases reported for Measles per epidemiological week



**Figure D3,** Total number of cases reported for Measles by each city for the entire time period




## Meningitis



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Meningitis (Meningococcus)

Indicator	Illinois	Chicago
Report period	1927-1964	1927-1948
Total weeks	1,280	1,015
Total cases	8,697	5,002
Max. cases per year	899	594
Year (max)	1944	1944
Max. cases per week	53	29
Week (max)	1944, wk 16	1944, wk 09
Average cases per year	334	227
95%CI	(238-430)	(154-300)
Average cases per week	7	5
95%CI	(7-7)	(5-5)



**Figure D2,** Number of cases reported for Meningitis per epidemiological week


## Meningococcal Disease



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Meningococcal Disease (Serogroup unspecified and Invasive all serogroups)

Indicator	Illinois	Chicago
Report period	1951-2009	1952-1953
Total weeks	2,608	80
Total cases	5,604	240
Max. cases per year	675	125
Year (max)	1953	1952
Max. cases per week	330	9
Week (max)	1952, wk 53	1953, wk 01
Average cases per year		
before 1980	118	120
95%CI	(60-176)	(56-184)
after 1980	75	-
95%CI	(62-88)	-
Average cases per week		
before 1980	3	3
95%CI	(2-4)	(3-3)
after 1980	2	-
95%CI	(2-2)	-



**Figure D2,** Number of cases reported for Meningococcal Disease per epidemiological week

## Mumps



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1,** Summary information for Mumps

Indicator	Illinois	Chicago	Springfield
Report period	1968-2002	1924-1932	1924-1932
Total weeks	1,651	450	417
Total cases	39,586	14,924	1,610
Max. cases per year	5,166	3,767	876
Year (max)	1968	1927	1925
Max. cases per week	439	193	63
Week (max)	1968, wk 08	1927, wk 13	1925, wk 10
Average cases per year			
before 1980	2,495	1,658	179
95%CI	(1,640-3,350)	(727-2,589)	(-30-388)
after 1980	325	-	-
95%CI	(4-646)	-	-
Average cases per week			
before 1980	49	33	4
95%CI	(45-53)	(29-37)	(3-5)
after 1980	7	-	-
95%CI	(6-8)	-	-





Figure D2, Number of cases reported for Mumps per epidemiological week


Pellagra



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Pellagra

Indicator	Chicago
Report period	1924-1932
Total weeks	436
Total cases	74
Max. cases per year	28
Year (max)	1928
Max. cases per week	2
Week (max)	1925, wk 38
Average cases per year	8
95%CI	(1-15)
Average cases per week	0
95%CI	(0-0)



**Figure D2,** Number of cases reported for Pellagra per epidemiological week

## Pneumonia



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1,** Summary information for Pneumonia (Unspecified)

Indicator	Illinois	Chicago
Report period	1948-1951	1912-1923
Total weeks	170	597
Total cases	12,783	119,212
Max. cases per year	4,778	28,396
Year (max)	1949	1918
Max. cases per week	364	7,743
Week (max)	1949, wk 13	1918, wk 41
Average cases per year	3,196	9,934
95%CI	(994-5,398)	(5,475-14,393)
Average cases per week	75	200
95%CI	(69-81)	(167-233)






Figure D2, Number of cases reported for Pneumonia per epidemiological week





Figure D3, Total number of cases reported for Pneumonia by each city for the entire time period

## Poliomyelitis



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1,** Summary information for Poliomyelitis (Total)

Indicator	Illinois	Chicago
Report period	1927-1970	1912-1953
Total weeks	1,861	1,531
Total cases	32,499	7,843
Max. cases per year	4,024	1,168
Year (max)	1952	1952
Max. cases per week	436	150
Week (max)	1952, wk 38	1943, wk 35
Average cases per year		
before 1960	951	201
95%CI	(620-1,282)	(117-285)
after 1960	19	-
95%CI	(1-37)	-
Average cases per week		
before 1960	20	5
95%CI	(18-22)	(4-6)
after 1960	1	-
95%CI	(1-1)	-




Figure D2, Number of cases reported for Poliomyelitis per epidemiological week





Figure D3, Total number of cases reported for Poliomyelitis by each city for the entire time period


## Rabies in Animals



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Rabies in Animals

Indicator	Illinois
Report period	1948-2009
Total weeks	2,451
Total cases	5,107
Max. cases per year	515
Year (max)	1980
Max. cases per week	52
Week (max)	1980, wk 23
Average cases per year	82
95%CI	(56-108)
Average cases per week	2
95%CI	(2-2)



**Figure D2,** Number of cases reported for Rabies in Animals per epidemiological week


## Rubella



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Rubella

Indicator	Illinois
Report period	1966-2001
Total weeks	1,515
Total cases	18,661
Max. cases per year	3,415
Year (max)	1968
Max. cases per week	997
Week (max)	1968, wk 11
Average cases per year	
before 1970	2,230
95%CI	(1,197-3,263)
after 1970	250
95%CI	(98-402)
Average cases per week	
before 1970	44
95%CI	(34-54)
after 1970	6
95%CI	(5-7)



**Figure D2,** Number of cases reported for Rubella per epidemiological week


## Salmonellosis



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1,** Summary information for Salmonellosis (PHLIS and Unspecified)


Indicator	Illinois
Report period	1999-2009
Total weeks	489
Total cases	12,510
Max. cases per year	1,742
Year (max)	2007
Max. cases per week	671
Week (max)	2004, wk 34
Average cases per year	1,137
95%CI	(786-1,488)
Average cases per week	26
95%CI	(22-30)



**Figure D2,** Number of cases reported for Salmonellosis per epidemiological week



## Scarlet Fever



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1,** Summary information for Scarlet Fever (Including streptococcal sore throat and Unspecified)

Indicator	Illinois	Chicago	Springfield
Report period	1927-1969	1906-1953	1907-1948
Total weeks	1,847	2,224	1,642
Total cases	395,035	274,461	6,066
Max. cases per year	35,046	17,025	601
Year (max)	1935	1935	1921
Max. cases per week	1,411	1,686	60
Week (max)	1963, wk 03	1907, wk 05	1920, wk 50
Average cases per year	10,677	6,099	148
95%CI	(8,176-13,178)	(5,052-7,146)	(107-189)
Average cases per week	214	123	4
95%CI	(204-224)	(118-128)	(4-4)





Figure D2, Number of cases reported for Scarlet Fever per epidemiological week



**Figure D3,** Total number of all cases reported for Scarlet Fever by each city for the entire time period


## Shigellosis



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Shigellosis (Unspecified and PHLIS)

Indicator	Illinois
Report period	1999-2009
Total weeks	489
Total cases	5,480
Max. cases per year	831
Year (max)	2003
Max. cases per week	283
Week (max)	2005, wk 50
Average cases per year	498
95%CI	(351-645)
Average cases per week	11
95%CI	(9-13)



**Figure D2,** Number of cases reported for Shigellosis per epidemiological week


## Streptococcal Disease, Invasive Group A



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1,** Summary information for Streptococcal Disease, Invasive Group A

Indicator	Illinois
Report period	2002-2009
Total weeks	373
Total cases	1,468
Max. cases per year	236
Year (max)	2009
Max. cases per week	71
Week (max)	2004, wk 24
Average cases per year	184
95%CI	(153-215)
Average cases per week	4
95%CI	(3-5)



**Figure D2,** Number of cases reported for Group A Streptococcal disease per epidemiological week


Streptococcal Sore Throat



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Streptococcal Sore Throat

Indicator	Illinois
Report period	1960-1961
Total weeks	102
Total cases	9,313
Max. cases per year	4,906
Year (max)	1961
Max. cases per week	324
Week (max)	1961, wk 10
Average cases per year	4,656
95%CI	(1,486-7,826)
Average cases per week	91
95%CI	(77-105)



**Figure D2,** Number of cases reported for Streptococcal Sore Throat per epidemiological week


## Streptococcus Pneumonia, Invasive Disease



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Streptococcus Pneumonia, Invasive Disease (Drug resistant <5 years)

Indicator	Illinois
Report period	2007-2008
Total weeks	77
Total cases	43
Max. cases per year	35
Year (max)	2007
Max. cases per week	26
Week (max)	2007, wk 45
Average cases per year	22
95%CI	(-150-194)
Average cases per week	1
95%CI	(0-2)



**Figure D2,** Number of cases reported for Streptococcus Pneumonia, Invasive Disease per epidemiological week


## Syphilis



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1,** Summary information for Syphilis (Primary and secondary and Civilian primary and secondary)


Indicator	Illinois
Report period	1972-2009
Total weeks	1,817
Total cases	27,399
Max. cases per year	2,739
Year (max)	1991
Max. cases per week	677
Week (max)	1991, wk 30
Average cases per year	721
95%CI	(527-915)
Average cases per week	15
95%CI	(14-16)



**Figure D2,** Number of cases reported for Syphilis per epidemiological week



## Tuberculosis



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1,** Summary information for Tuberculosis (Unspecified)

Indicator	Illinois	Chicago
Report period	1975-2005	1906-1923
Total weeks	1,485	857
Total cases	32,806	157,949
Max. cases per year	1,733	16,609
Year (max)	1977	1918
Max. cases per week	413	547
Week (max)	1991, wk 30	1917, wk 50
Average cases per year	1,058	8,775
95%CI	(908-1,208)	(6,362-11,188)
Average cases per week	22	184
95%CI	(21-23)	(177-191)






Figure D2, Number of cases reported for Tuberculosis per epidemiological week



**Figure D3,** Total number of cases reported for Tuberculosis by each city for the entire time period


## Typhoid Fever




**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1,** Summary information for Typhoid Fever (Unspecified and Including paratyphoid fever)

Indicator	Illinois	Chicago	Springfield
Report period	1927-2005	1906-1953	1907-1948
Total weeks	2,984	2,074	1,371
Total cases	13,441	16,004	762
Max. cases per year	1,167	2,020	70
Year (max)	1934	1909	1915
Max. cases per week	259	264	14
Week (max)	1939, wk 35	1909, wk 09	1917, wk 36
Average cases per year			
before 1950	505	372	20
95%CI	(339-671)	(218-526)	(14-26)
after 1950	28	12	-
95%CI	(13-43)	(-32-56)	-
Average cases per week			
before 1950	10	8	1
95%CI	(9-11)	(7-9)	(1-1)
after 1950	1	1	-
95%CI	(1-1)	(1-1)	-




**Figure D2,** Number of cases reported for Typhoid Fever per epidemiological week



**Figure D3,** Total number of all cases reported for Typhoid Fever by each city for the entire time period

## Whooping Cough



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1,** Summary information for Whooping Cough

Indicator	Illinois	Chicago	Springfield	Elgin
Report period	1937-2009	1906-1953	1908-1948	1935-1941
Total weeks	2,268	1,541	1,094	334
Total cases	100,120	105,292	3,425	894
Max. cases per year	14,325	9,951	392	210
Year (max)	1938	1938	1934	1936
Max. cases per week	616	422	68	26
Week (max)	2005, wk 50	1938, wk 48	1943, wk 03	1936, wk 47
Average cases per year				
before 1960	4,938	3,290	132	128
95%CI	(2,920-6,956)	(2,506-4,074)	(88-176)	(65-191)
after 1960	180	-	-	-
95%CI	(97-263)	-	-	-
Average cases per week				
before 1960	102	68	3	3
95%CI	(96-108)	(65-71)	(3-3)	(3-3)
after 1960	5	-	-	-
95%CI	(4-6)	-	-	-




Figure D2, Number of cases reported for Whooping Cough per epidemiological week



## Diseases with fragmented data over time

For a number of diseases, only fragmented data was available for Illinois. This was due to the inclusion of these diseases in the weekly surveillance system for short periods or widely dispersed periods of time. Due to these wide gaps or limited data, data for these diseases were not presented in separate sections but in the table below. Work will continue to include more data in the Tycho database and complete data sets for each disease where possible. Collaborations with local, state and federal public health agencies will be made to explore data availability and opportunities to include these in the Tycho database.

**Table 2,** Summary information on the occurrence of diseases with fragmented data in Illinois

Disease	Report type	Report period	Number of reports	Total cases
Anthrax	State	1942-1945	143	1
Botulism	City	1952-1952	1	1
Brucellosis [undulant fever]	City	1952-1981	27	41
Brucellosis [undulant fever]	State	1943-1982	756	6187
Dengue	City	1924-1925	9	0
Dysentery	State	1942-1948	326	803
Encephalitis	City	1941-1981	635	67
Encephalitis	State	1942-1994	1608	2972
Leprosy	City	1924-1924	1	0
Leprosy	State	1942-1990	387	22
Listeriosis	State	2000-2005	133	45
Lyme disease	State	1991-2009	424	458
Psittacosis	State	1956-1961	48	64
Rocky mountain spotted fever	City	1952-1953	3	18
Rocky mountain spotted fever	State	1942-2009	1219	656
Tetanus	State	1962-1980	489	104
Toxic shock syndrome	State	1983-1994	265	97
Trichiniasis	City	1952-1953	4	6
Trichiniasis	State	1952-1955	8	12
Tularemia	City	1952-1953	4	6
Tularemia	State	1942-1994	1060	1114
Typhus fever	City	1924-1925	48	0
Typhus fever	State	1942-1958	200	21
West nile disease	State	2003-2009	127	337

## Project Tycho

This report provides preliminary data for the state of Illinois available in the Tycho database. This database is currently being beta tested and these data cannot be used for publication or other official use at this time. An open access release to the general public is planned for later in 2011.

Please visit the Tycho website for more information and to query the database at: [www.tycho.pitt.edu](http://www.tycho.pitt.edu). For further information regarding the Tycho project, contact Dr. Wilbert van Panhuis at the University of Pittsburgh Graduate School of Public Health.

Dr. Wilbert van Panhuis, MD PdD  
Graduate School of Public Health  
University of Pittsburgh  
130 DeSoto Street  
704 Parran Hall  
Pittsburgh, 15261 PA  
Tel: 412-624-7693  
Email: [wav10@pitt.edu](mailto:wav10@pitt.edu)