

Preliminary data, not for publication or official use

# Project Tycho

***Preliminary data for the state of New Mexico***

## Tycho database beta test version

*The data presented in this report are of preliminary nature and should not be used for publication or other types of official use*

June 2011

University of Pittsburgh Graduate School of Public Health



University of Pittsburgh



## Table of Content

Acknowledgements	page 3
Contact information	page 4
Introduction	page 5
Vision and goal	page 7
Collaboration	page 8
Summary of methods	page 9
Overview of data available for the state of New Mexico	page 12
Disease specific data for New Mexico	page 18
- AIDS	page 18
- Chickenpox	page 19
- Chlamydia	page 20
- Coccidioidomycosis	page 21
- Cryptosporidiosis	page 22
- Diphtheria	page 23
- Escherichia Coli	page 25
- Giardiasis	page 26
- Gonorrhea	page 27
- Haemophilus Influenzae	page 28
- Hepatitis	page 29
- Legionellosis	page 31
- Listeriosis	page 32
- Measles	page 33
- Meningitis	page 35
- Meningococcal disease	page 36
- Mumps	page 38
- Pneumonia	page 39
- Poliomyelitis	page 40
- Rabies in animals	page 42
- Salmonellosis	page 43
- Scarlet fever	page 45
- Shigellosis	page 46
- Streptococcal disease, invasive group A	page 47
- Streptococcal sore throat	page 48
- Syphilis	page 49
- Tuberculosis	page 51
- Typhoid fever	page 51
- Typhus fever	page 53

# Preliminary data, not for publication or official use

## Disease specific data for New Mexico, continued

- Whooping cough \_\_\_\_\_ **page 54**

**Diseases with fragmented data** \_\_\_\_\_ **page 56**

## Acknowledgements

The development of this project would not have been possible without data collected every week by staff from city-, county-, state- and federal public health departments. We would like to acknowledge their invaluable contribution to the overall body of surveillance data that were digitized during this project. Particularly, we acknowledge staff from the Council of State and Territorial Epidemiologists (CSTE) and the US Centers for Disease Control for their overall efforts in disease surveillance and their particular input in this project.

We also acknowledge the help of many contributors that have contributed to the digitization and processing of weekly surveillance data included in this database. Digital Divide Data: Linda Thomas, Eric Gold, Heak Hok, Villa Kem and Hetel Patel. University of Pittsburgh Graduate School of Public Health: Steve Wisniewsky, Heather Eng, Sharon Crow, David Galloway, Jia Xu and Raaka Kumbhakar. University of Pittsburgh School of Information Sciences: Vladimir Zadorozhny, Ying-Feng Hsu, David Paul Koenig, Fatimah Ridwan and Divyasheel Sharma. Johns Hopkins Bloomberg School of Public Health: Derek Cummings. Pittsburgh Supercomputing Center: David Graham and Andrew Adams.

This project was made possible with funding from the Bill & Melinda Gates Foundation through the Vaccine Modeling Initiative (grant ID 49276).

The Tycho development team at the University of Pittsburgh Graduate School of Public Health:

Willem G. van Panhuis

John Grefenstette

Shawn T. Brown

Su Yon Jung

Nian Shong Chok

Anne Cross

Donald S. Burke

## Contact information

The Tycho database is currently being tested and a login account will be required to use the database and website ([www.tycho.pitt.edu](http://www.tycho.pitt.edu)). All State Epidemiologists, CDC partners, other federal agencies and our research collaborators will be provided with user accounts. User accounts can also be provided upon specific request. Note that the quality of the data during the testing phase cannot be guaranteed to be sufficient for publication or official use.

**Point of contact for the Tycho project:**

Dr. Willem G. van Panhuis, MD PhD

Graduate School of Public Health

University of Pittsburgh

130 DeSoto Street

704 Parran Hall

Pittsburgh PA, 15261

Email: [wav10@pitt.edu](mailto:wav10@pitt.edu)

Tel: 412-624-7693

## Introduction

**Project Tycho** started in 2009 as part of the Vaccine Modeling Initiative (VMI) at the University of Pittsburgh Graduate School of Public Health, funded by the Bill & Melinda Gates foundation. This project aims to contribute to the availability of public health data for research and policy making. The vision for this project as described in the next section is a central access point for public health data of a detailed spatial and temporal scale for all countries. Currently, the Tycho database contains 10 million records that each represent a weekly report from a location for a specific disease. These records were extracted from weekly US Nationally Notifiable Disease Surveillance Reports between 1888 and 2009 (6300 reports) using double data entry (200 million keystrokes). The current database includes a total of 90 million reported cases and 4 million reported deaths due to notifiable diseases in the United States for the last 122 years.

Project Tycho is named after the Danish nobleman Tycho Brahe (1546 – 1601), who meticulously collected astronomical data. After Brahe's death, his assistant Kepler used these data to develop area laws of planetary motion.

## Background

***"No health department, State or local, can effectively prevent or control diseases without knowledge of when, where and under what conditions cases are occurring"***

(Quote stated above weekly surveillance reports published in the Public Health Reports, 1888-1951)

This quote has accompanied weekly reports on the prevalence of notifiable diseases in the United States as long as these were published in the Public Health Reports, emphasizing the importance of data for disease control policy making. In the current day and age, this statement has lost none of its significance. On the contrary, as infectious diseases continue to pose challenges to the global public health system in an increasingly interconnected global society. The public health system has responded to this challenge by evermore sophisticated disease surveillance systems. The availability of internet and information technology has facilitated collection of detailed data on infectious disease incidence, the spread of pathogens, disease determinants, health behavior, etc. New technologies such as remote sensing and mobile reporting systems have advanced measurement of these factors in both high and low resource settings. The enormous wealth of public health information that is currently being accrued also poses its own challenges for data storage, management, preservation and dissemination. Public health agencies will need to allocate resources and develop capacity to specifically address these issues.

Although the importance of public health data collection has been widely recognized, the dissemination of data has been neglected for many reasons. The above quote does not only imply data collection but also its use and dissemination as knowledge on disease occurrence can only be derived from data after processing and analysis. Not only has disease surveillance become more sophisticated, analytical tools have changed as well and now include computational models to assess disease transmission and to evaluate disease control options. These methods have introduced the opportunity for data intensive studies of fine spatial-temporal patterns using dynamic models. Disease surveillance data at such scale are often not available for analysis for multiple reasons. First, surveillance data are not collected for research purposes but for disease monitoring and planning. Secondly, public health staff often lack time or capacity to prepare these data for analysis. Third, surveillance methodology often lacks transparency and standardization, reducing options for scientific analyses. Fourth, disease surveillance data are often

# Preliminary data, not for publication or official use

kept confidential and ownership or data use requirements are often unclear. Finally, archiving and preservation of surveillance data are often not a priority and disaggregated records may get lost, losing the opportunity to ever use these data for analysis.

The Tycho project aims at overcoming some of these challenges by providing open access to large public health datasets of great spatial-temporal detail. Open access to these data will provide great opportunities for scientific analysis and better informed policy making on disease control.

## **The Tycho database and website**

As described in the methods section, a large heterogeneity in reporting practice was observed over the past 122 years of weekly disease surveillance and substantial efforts were made to clean and standardize these data. The current online database includes all cleaned and standardized data in a format that allows comparisons across time and space. It does not include parts of the data that require additional cleaning and research such as classification of disease subcategories.

Each record in the database provides a reported number of cases or deaths due to a specific disease for a specific location, for a certain week. The database can be searched online ([www.tycho.pitt.edu](http://www.tycho.pitt.edu)) by selecting a dataset, an aggregation method and an output format. Data can be viewed in tables, graphics and maps and these can be downloaded for further use.

## **Data availability**

The data provided by the Tycho database reflect the weekly US Nationally Notifiable Disease Surveillance System between 1888 and 2009. It includes all diseases that were reported as part of this system, but it does not yet include records from other surveillance systems such as the annual system or disease specific subsystem (such as the influenza surveillance system).

The available data also reflects changes in reporting practices over time. This explains why city or state level reports are only available for a certain time period (1888-1953 for cities and 1927-2009 for states) and why morbidity and mortality reports are available for different time periods. We are currently working on inclusion of additional data as described in the section on collaboration.

## **Testing and release of the Tycho database**

The Tycho database is currently in a beta testing phase that will start during the 2011 annual meeting of the Council of State and Territorial Epidemiologists (CSTE) in Pittsburgh June 12-16. This phase will end with a release to the general public in the fall of 2011. During the testing phase, invited users will be provided with login accounts that will enable full use of all features of the website and database. Invited users will include all state epidemiologists, partners in the CDC and other federal agencies as well as research collaborators. During this phase, data cleaning and standardization will continue. The quality of the data will not be sufficient for publication or official use during this phase.

## **Future developments**

After optimization of the database and website during the testing phase, the first version of the Tycho database will be released to the general public. After that, efforts to clean and standardize the data will continue and more data will be released in subsequent versions. In addition, ongoing and new collaborations for inclusion of new data will allow gradual expansion of the dataset over time and increased opportunities for data driven analysis and policy making.

## Vision and goal

The vision for project Tycho is based on the value of open access to detailed, disaggregated public health data for scientific analysis and policy making. Starting with 122 years of weekly US surveillance data, it is our hope that this example will convince public health authorities worldwide of the value of this resource and will stimulate similar contributions from other countries.

In the past, data sharing and archiving have not been a priority in public health or among health sciences in general. Continuous and diligent public health data collection has led to a large pool of disaggregated data scattered over the world including the internet, ranging from paper archives to basements of local health departments. The need for open access to a central repository of public health data has been recognized and both benefits and challenges of open access to data in general have been well described. Open access to data could lead to increased accountability, transparency, innovation, collaboration, cost-effectiveness, replication of results, the development of new methods and insights, and a narrower data access gap between low- and higher income countries. Ultimately, this would result in the advancement of science and technology to the benefit of all.

Challenges to open access include inconsistent formats, lack of annotation or metadata, lack of information technology (IT) capacity, a lack of incentive, and cultural norms around data ownership. Open access to public health data in particular is confronted by decentralized archiving of disease reports, lack of time and capacity for data management, degraded physical format of records and archives, and governmental level legal constraints. Multiple principles and requirements for a central open access repository for public health data have been described and include (1) capacity building for data management in low- and middle income countries, (2) feedback loops to data contributors, (3) sustainability, (4) common standards, (5) interoperability, and (6) user-friendliness.

The ultimate goal of the Tycho project is to provide a central global public health data access point. Historical as well as current public health data are of great value if archived and accessible for research and analysis. Open access will enable the use of analytical capacity from around the globe which will lead to new discoveries of disease patterns and control policies. Furthermore, a central data access point will facilitate data archiving and preservation into the future which will be an increasing need in a data rich public health environment.

## Collaboration

The application of the surveillance data from the Tycho database as well as the inclusion of new data will require a wide range of collaborations with public health offices in the US and abroad. The Tycho project has currently made the first step of digitizing 122 years of weekly US surveillance reports that had already been published. The next step will be the completion of the existing dataset as well as the inclusion of new data from unpublished sources.

This report provides an overview of data available from the Tycho database for the state of New Mexico. This includes all data that has ever been published at state or city level for New Mexico in the weekly US Nationally Notifiable Disease Surveillance System. As you will find, the data availability will vary greatly over time, between city and state level reports, between diseases and between morbidity vs. mortality reports.

The Tycho development team would look forward to work with state health departments in the US to:

1. Better understand the current Tycho data for each state (eg. by collecting historic documentation)
2. Provide better usability and applications of current Tycho data for each state by continued joint development of website and database features.
3. Collect and/or digitize official, confirmed data to validate the current (preliminary) weekly data.
4. For each state, collect and/or digitize new data that has not yet been included. For example all city level reports discontinued in 1953 and completion of city level data until 2009 would greatly increase opportunities for analysis of disease patterns and trends.
5. Provide support to state health offices to manage, preserve and provide access to public health data.

## Summary of methods

Detailed documentation on the methods used for the Tycho database has been provided on the website ([www.tycho.pitt.edu](http://www.tycho.pitt.edu)). This section describes these methods in short.

### Data collection

Weekly reports that contain tables on the occurrence of nationally notifiable diseases have been published since 1888 by public health authorities at the Federal level in various journals. Table 1 provides the list of publications and the responsible agency since 1888. All weekly nationally notifiable disease reports between 1888 and 1951 could be retrieved from the PubMed Central repository of the National Library of Medicine <sup>1</sup>. For 1995 to 2009, these reports could be retrieved from the MMWR digital archive on the CDC website <sup>2</sup>. Most weekly reports between 1952 and 1995 could be retrieved from the HathiTrust Digital Library <sup>3 4</sup>, but many could not be found and had to be copied from hard copies of MMWR issues in the University of Pittsburgh library.

Table 1, publications and responsible Federal agencies for nationally notifiable disease reports

Time period	Publication title	Responsible federal agency
1888-1889	Weekly Abstract of Sanitary Reports	US Marine Hospital Service
1890-1895	Abstract of Sanitary Reports	US Marine Hospital Service
1896-1901	Public Health Reports	US Marine Hospital Service
1902-1911	Public Health Reports	US Public Health and Marine Hospital Service
1912-1951	Public Health Reports	US Public Health Service
1952-1960	Morbidity and Mortality Weekly Report	National Office of Vital Statistics, US Public Health Service
1961-1969	Morbidity and Mortality Weekly Report	Communicable Disease Center
1970-1991	Morbidity and Mortality Weekly Report	Center for Disease Control
1992-2009	Morbidity and Mortality Weekly Report	Centers for Disease Control and Prevention

### Inclusion criteria

Weekly reports of each year were reviewed systematically to assess the diseases reported. We included all tables that provided disease specific information by week for US cities, townships, counties or states. Tables that provided summary or aggregated information by month, year or at the national level were not included. Similarly, tables that did not contain disease specific information (such as all cause mortality) were not included.

### Data entry

Weekly reports were downloaded or scanned as PDF files and selected tables with notifiable disease reports were entered into computer spreadsheets in a highly standardized fashion using double data entry. During the second round of data entry, operators could not see what had been entered in the first round and could not continue if the system detected a discrepancy between the second and first entry

<sup>1</sup> <http://www.ncbi.nlm.nih.gov/pmc/tocrender.fcgi?journal=333&action=archive>

<sup>2</sup> <http://www.cdc.gov/mmwr>

<sup>3</sup> <http://catalog.hathitrust.org/Record/003910026>

<sup>4</sup> <http://catalog.hathitrust.org/Record/003843660>

# Preliminary data, not for publication or official use

for a specific value. Such discrepancies could only be resolved by checking the PDF file and try again or discussion with the group leader and verification of a value in the source documents.

## **Quality control for data entry**

The accuracy of data entry was checked at various levels. First, completeness of data was verified by comparing the content of entered data with PDF sources files. Secondly, accuracy of data entry was verified by multiple rounds of comparing random samples of entered files with PDF source files. Thirdly, data formatting was verified by various checks to ensure appropriate formatting for data loading.

## **Data loading and standardization**

All data was entered in Excel spreadsheets and various components of these spreadsheets were loaded in data files. Table titles, column headers, place names and reported numbers were loaded in separate files. These files were used to extract information on each reported number, including:

1. the disease reported
2. the disease subcategory reported
3. cases or deaths reported
4. the reporting location (name, state and type of location)
5. the time period for which a number was reported
6. the date of publication of the original weekly report associated with a reported number

## **Integration**

All reported numbers and extracted information was integrated in one database with one record per reported number and associated information.

## **Post-processing quality control**

After integration of all data in one database, checks were performed to detect duplicate reports and data inconsistencies. Duplicate records were removed and inconsistencies resolved by verification with original PDF source files.


## **Data filtering**

The digitized version of all historical weekly US nationally notifiable disease surveillance records is a very heterogeneous dataset. It took a substantial number of data processing protocols to standardize time and space variables as well as disease names. All standardized records have been separated from non-standard records by a filter. All remaining heterogeneity in non-standard records is inherent to the surveillance system and can only be standardized after further analysis will have been completed (eg. remaining heterogeneity in reporting periods for reports before 1953 and standardization of disease subcategories). The largest proportion of data (>4 million records) has been standardized however and has been made available in the current testing version.

## **Data visualization**

Figures in this report were made with the R system, version 2.9.2 and the maps were generated by the GAIA platform developed at the University of Pittsburgh Graduate School of Public Health in collaboration with the Pittsburgh Supercomputing Center (PSC). See <http://midas-pitt.psc.edu/gaia> for more information

# Preliminary data, not for publication or official use



**Figure 1, Schematic of the Tycho data architecture.** Access to a consistent, standardized subset of the digitized US weekly surveillance reports will be provided for beta testing through a website that will allow querying and downloading of data. Data from the development database will be pushed through a filter to feed the Tycho database.

The current Tycho database will provide the reported number of cases or deaths reported by city or state health authorities to the federal health agency for all weeks between 1888 and 2009. Currently, only standardized, consistent data are being released for testing, according to the following criteria:

1. Only weekly reports are provided (reports for other periods such as 10 days, 2 weeks, 1 month, 1 year) are currently not included due to lack of comparability with the weekly reports.
2. Only reports that were published in the same year as the reporting period were included (this excludes updates or comparisons that were published more than a year after the original reporting period)
3. Only numbers were included for which information about the location, time period and disease could be extracted.
4. Only numbers or time periods for which no disease subcategories were reported were included.

Each of the steps described above have been described in detailed protocols that are posted in the documentation part of the Tycho website ([www.tycho.pitt.edu](http://www.tycho.pitt.edu)).

## Website

The Tycho website features simple and advanced searches of the data, visualizations and downloading of machine readable files. A dataset can be selected by specifying: 1) the disease of interest, 2) the location(s) of interest and 3) the time period. In a second step, the type of aggregation and length of increments can be specified and in a third step, the type of output (table, graphic or map) can be selected. After output has been generated, data can be downloaded.

As mentioned above, data availability varies largely by disease, time period and location. We aimed to restrict selection options by data availability to avoid "no results available."

Access to the most of the website will be password protected during the testing phase. For invited users after login, full functionality of the database and website will be available. During this phase however, the data quality will not be sufficient for publication or official use.

## Overview of data available for the state of New Mexico

In this section, an overview of data available for the state of New Mexico will be provided. As described earlier, data availability depends entirely on historical reporting practices in the weekly US Nationally Notifiable Disease Surveillance System between 1888 and 2009. For this report, we only included morbidity reports (cases) to improve the format and limit the size of this report. As mentioned earlier, the analysis and standardization of has not been completed yet for all records in the database and preliminary data are provided here that may not yet be available in the online database.

This section will provide a general overview of data availability. The next sections will provide disease specific data from state and city level reports of New Mexico. Data for a maximum of 3 major cities were provided depending on availability (city data was only reported until 1953). Some summary data will be provided at the end of this report, for a subset of diseases for which data was fragmented over time.

Table 1 lists the number of weekly state or city reports that are available for New Mexico per each disease and subcategory.

**Table 1, Number of weekly state or city reports per disease and subcategory**

Disease	City	State
Aids	-	912
Anthrax	-	143
Botulism	-	1
Brucellosis [undulant fever]	1	145
Chickenpox [varicella]	425	532
Chlamydia	-	634
Coccidioidomycosis	-	280
Cryptosporidiosis	-	435
Dengue	2	-
Diphtheria	982	1722
Dysentery		
<i>Amebic</i>	-	222
<i>Bacillary</i>	-	252
<i>Unspecified</i>	-	240
Encephalitis		
<i>Lethargic</i>	62	-
<i>Post infectious</i>	-	98
<i>Primary [infectious] including unspecified</i>	1	635
Escherichia coli		
<i>EHEC 0157</i>	-	143
<i>EHEC non-0157</i>	-	132
<i>O157:H7 NETSS</i>	-	248
<i>O157:H7 PHLIS</i>	-	187
<i>STEC</i>	-	177
Giardiasis	-	367

# Preliminary data, not for publication or official use

**Table 1.** Number of weekly state or city reports per disease and subcategory, *continued*

Disease	City	State
Gonorrhea		
<i>Civilian</i>	-	755
<i>Unspecified</i>	-	1047
Haemophilus influenzae		
<i>Age &lt;5 non-serotype B</i>	-	174
<i>Age &lt;5 serotype B</i>	-	13
<i>Age &lt;5 unknown serotype</i>	-	132
<i>All ages all serotypes</i>	-	633
Hepatitis		
<i>Acute type A</i>	-	352
<i>Acute type B</i>	-	353
<i>Acute type C</i>	-	34
<i>Acute type NA NB [including C]</i>	-	19
<i>All types, &lt;20 years</i>	-	191
<i>All types, &gt;=20 years</i>	-	76
<i>All types, all ages</i>	-	645
<i>Type A [infectious]</i>	-	1658
<i>Type B [serum]</i>	-	1111
<i>Type NA NB [including C]</i>	-	727
<i>Type unspecified</i>	-	531
Influenza	334	734
Legionellosis	-	823
Leprosy	-	168
Listeriosis	-	155
Lyme disease	-	382
Malaria	-	1353
Measles		
<i>Imported</i>	-	312
<i>Indigenous</i>	-	394
<i>Unspecified</i>	977	2423
Meningitis		
<i>Aseptic</i>	-	569
<i>Meningococcus</i>	90	1128
<i>Unspecified</i>	15	5
Meningococcal disease		
<i>All serogroups</i>	-	34
<i>Invasive all serogroups</i>	-	163
<i>Invasive serogroup unknown</i>	-	14
<i>Serogroup unspecified</i>	10	1943
Mumps	423	707
Pellagra	62	-

# Preliminary data, not for publication or official use

**Table 1.** Number of weekly state or city reports per disease and subcategory, *continued*

Disease	City	State
Pneumonia	17	166
Poliomyelitis		
<i>Non paralytic</i>	-	71
<i>Paralytic</i>	-	193
<i>Total</i>	136	1675
Psittacosis	-	2
Rabies in animals	-	1944
Rocky mountain spotted fever	1	674
Rubella	-	781
Salmonellosis		
<i>NETSS</i>	-	127
<i>PHLIS</i>	-	123
<i>Unspecified</i>	-	365
Scarlet fever		
<i>Including streptococcal sore throat</i>	48	544
<i>Unspecified</i>	972	1238
Shigellosis		
<i>NETSS</i>	-	126
<i>PHLIS</i>	-	123
<i>Unspecified</i>	-	374
Streptococcal disease, invasive group a	-	374
Streptococcal sore throat	-	99
Streptococcus pneumoniae invasive disease		
<i>Drug resistant A</i>	-	5
<i>Drug resistant all ages</i>	-	195
<i>Non drug resistant &lt;5 years</i>	-	133
Syphilis		
<i>Civilian primary and secondary</i>	-	697
<i>Congenital</i>	-	82
<i>Primary and secondary</i>	-	925
Tetanus	-	54
Toxic shock syndrome	-	200
Trichiniasis	1	-
Tuberculosis [phthisis pulmonalis]		
<i>New active</i>	-	135
<i>Unspecified</i>	116	1400
Tularemia	-	855
Typhoid fever [enteric fever]		
<i>Including paratyphoid fever</i>	-	638
<i>Unspecified</i>	949	1869

# Preliminary data, not for publication or official use


**Table 1.** Number of weekly state or city reports per disease and subcategory, *continued*

Disease	City	State
Typhus fever		
<i>Endemic</i>	-	22
<i>Unspecified</i>	6	175
West nile disease		
<i>Neuroinvasive</i>	-	140
<i>Non-neuroinvasive</i>	-	95
Whooping cough [pertussis]	893	2233

The diseases included in the weekly US Nationally Notifiable Disease Surveillance System varied largely over time and reflected the historical social-political priorities of each time period. Note that the diseases in the weekly system were a subset of all diseases included in the annual Notifiable Disease Surveillance System (that was not entered as part of this project).

The number of locations reporting diseases also varied over time depending on the type of reports submitted. Figure 1 on the next page provides an overview of the weeks for which a report was available per disease at the state or city level. Colors indicate the total number of reporting locations.


# Preliminary data, not for publication or official use




**Figure 1,** The number of locations (in color code) reporting on each disease per week between 1888 and 2009

The map in figure 2 (next page) illustrates the geographical distribution of the available city data. It displays the number of weekly reports for any disease available per location that was included (and for which coordinates could be derived)

Figure 3 below provides an overview of the number of diseases for which a weekly report was included at the state level and for individual cities. All cities for which at least 100 weekly reports were available have been listed. For most states, it shows a major drop in the number of cities after 1925 and after 1953, city reports were discontinued except for New York City, which became its own reporting jurisdiction.




**Figure 2,** This map displays the number of weekly reports available for any disease per city (note that city reports were only available until 1953)



**Figure 3,** The number of diseases (in color code) that were reported for the state and city level (state on top row) for each week between 1888 and 2009

## Disease specific data for New Mexico


### AIDS



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1, Summary information for AIDS**

Indicator	New Mexico
Report period	1984-2005
Total weeks	912
Total cases	2,493
Max. cases per year	282
Year (max)	1993
Max. cases per week	155
Week (max)	2004, wk 22
Average cases per year	113
95%CI	(79-147)
Average cases per week	3
95%CI	(2-4)



**Figure D2,** Number of cases reported for AIDS per epidemiological week


## Chickenpox



**Figure D1**, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1**, Summary information for Chickenpox

Indicator	Albuquerque
Report period	1924-1932
Total weeks	425
Total cases	1,101
Max. cases per year	263
Year (max)	1931
Max. cases per week	19
Week (max)	1930, wk 49
Average cases per year	
<i>before 1995</i>	122
95%CI	(74-170)
Average cases per week	
<i>before 1995</i>	3
95%CI	(3-3)



**Figure D2**, Number of cases reported for Chickenpox per epidemiological week


## Chlamydia



**Figure D1**, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1**, Summary information for Chlamydia

Indicator	New Mexico
Report period	1996-2009
Total weeks	634
Total cases	73,489
Max. cases per year	9,391
Year (max)	2004
Max. cases per week	3,667
Week (max)	2004, wk 22
Average cases per year	5,249
95%CI	(4,025-6,473)
Average cases per week	116
95%CI	(97-135)



**Figure D2**, Number of cases reported for Chlamydia per epidemiological week


## Coccidioidomycosis



**Figure D1**, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1**, Summary information for Coccidioidomycosis

Indicator	New Mexico
Report period	2003-2009
Total weeks	280
Total cases	1,219
Max. cases per year	1,129
Year (max)	2004
Max. cases per week	1,109
Week (max)	2004, wk 22
Average cases per year	174
95%CI	(-215-563)
Average cases per week	4
95%CI	(-4-12)



**Figure D2**, Number of cases reported for Coccidioidomycosis per epidemiological week


## Cryptosporidiosis



**Figure D1**, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1**, Summary information for Cryptosporidiosis

Indicator	New Mexico
Report period	1999-2009
Total weeks	435
Total cases	462
Max. cases per year	113
Year (max)	2009
Max. cases per week	16
Week (max)	2008, wk 35
Average cases per year	42
95%CI	(16-68)
Average cases per week	1
95%CI	(1-1)



**Figure D2**, Number of cases reported for Cryptosporidiosis per epidemiological week


## Diphtheria



**Figure D1**, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1**, Summary information for Diphtheria

Indicator	New Mexico	Albuquerque
Report period	1927-1977	1918-1941
Total weeks	1,722	982
Total cases	5,277	578
Max. cases per year	591	121
Year (max)	1932	1922
Max. cases per week	366	9
Week (max)	1962, wk 43	1922, wk 28
Average cases per year		
<i>before 1940</i>	255	26
95%CI	(167-343)	(14-38)
<i>after 1940</i>	50	12
95%CI	(21-79)	-
Average cases per week		
<i>before 1940</i>	6	1
95%CI	(6-6)	(1-1)
<i>after 1940</i>	2	0
95%CI	(1-3)	(0-0)



**Figure D2,** Number of cases reported for Diphtheria per epidemiological week


## Escherichia Coli



**Figure D1**, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1**, Summary information for Escherichia Coli (O157:H7 PHLIS, EHEC 0157, and STEC)

Indicator	New Mexico
Report period	1996-2009
Total weeks	507
Total cases	214
Max. cases per year	45
Year (max)	2008
Max. cases per week	7
Week (max)	1998, wk 48
Average cases per year	15
95%CI	(7-23)
Average cases per week	0
95%CI	(0-0)



**Figure D2**, Number of cases reported for Escherichia Coli per epidemiological week


## Giardiasis




**Figure D1**, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1, Summary information for Giardiasis**

Indicator	New Mexico
Report period	2002-2009
Total weeks	367
Total cases	6,475
Max. cases per year	5,865
Year (max)	2006
Max. cases per week	5,801
Week (max)	2006, wk 35
Average cases per year	809
95%CI	(-899-2,517)
Average cases per week	18
95%CI	(-13-49)




## Gonorrhea



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Gonorrhea (Unspecified and Civilian)

Indicator	New Mexico
Report period	1972-2009
Total weeks	1,802
Total cases	92,292
Max. cases per year	7,080
Year (max)	1975
Max. cases per week	1,191
Week (max)	2006, wk 36
Average cases per year	2,429
95%CI	(1,805-3,053)
Average cases per week	51
95%CI	(48-54)



**Figure D2,** Number of cases reported for Gonorrhea per epidemiological week


## Haemophilus Influenzae



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Haemophilus Influenzae (All ages all serotypes)

Indicator	New Mexico
Report period	1996-2009
Total weeks	633
Total cases	341
Max. cases per year	39
Year (max)	2007
Max. cases per week	11
Week (max)	2005, wk 26
Average cases per year	24
95%CI	(18-30)
Average cases per week	1
95%CI	(1-1)



**Figure D2,** Number of cases reported for Haemophilus Influenzae per epidemiological week


## Hepatitis



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Hepatitis (Type A [Infectious], Acute type A, Type B [Serum], and Acute type B)

Indicator	New Mexico
Report period	1966-2009
Total weeks	2,042
Total cases	12,464
Max. cases per year	737
Year (max)	1994
Max. cases per week	376
Week (max)	1970, wk 44
Average cases per year	
<i>before 1990</i>	357
<i>95%CI</i>	(303-411)
<i>after 1990</i>	187
<i>95%CI</i>	(86-288)
Average cases per week	
<i>before 1990</i>	8
<i>95%CI</i>	(7-9)
<i>after 1990</i>	4
<i>95%CI</i>	(4-4)



**Figure D2,** Number of cases reported for Hepatitis per epidemiological week


## Legionellosis



**Figure D1**, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1**, Summary information for Legionellosis

Indicator	New Mexico
Report period	1982-2009
Total weeks	823
Total cases	95
Max. cases per year	9
Year (max)	2007
Max. cases per week	2
Week (max)	1983, wk 01
Average cases per year	3
95%CI	(2-4)
Average cases per week	0
95%CI	(0-0)



**Figure D2**, Number of cases reported for Legionellosis per epidemiological week


## Listeriosis



**Figure D1**, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1, Summary information for Listeriosis**

Indicator	New Mexico
Report period	2000-2005
Total weeks	155
Total cases	13
Max. cases per year	6
Year (max)	2001
Max. cases per week	2
Week (max)	2001, wk 25
Average cases per year	2
95%CI	(0-4)
Average cases per week	0
95%CI	(0-0)



**Figure D2**, Number of cases reported for Listeriosis per epidemiological week


## Measles



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1,** Summary information for Measles (Unspecified)

Indicator	New Mexico	Albuquerque
Report period	1927-1997	1918-1953
Total weeks	2,423	977
Total cases	82,567	4,751
Max. cases per year	8,082	626
Year (max)	1958	1932
Max. cases per week	1,313	101
Week (max)	1953, wk 17	1927, wk 07
Average cases per year		
<i>before 1970</i>	1,940	183
95%CI	(1,358-2,522)	(101-265)
<i>after 1970</i>	79	-
95%CI	(19-139)	-
Average cases per week		
<i>before 1970</i>	43	5
95%CI	(40-46)	(4-6)
<i>after 1970</i>	2	-
95%CI	(1-3)	-



**Figure D2,** Number of cases reported for Measles per epidemiological week


## Meningitis



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Meningitis

Indicator	New Mexico
Report period	1927-1964
Total weeks	1,128
Total cases	583
Max. cases per year	80
Year (max)	1930
Max. cases per week	19
Week (max)	1940, wk 20
Average cases per year	22
95%CI	(14-30)
Average cases per week	1
95%CI	(1-1)



**Figure D2,** Number of cases reported for Meningitis per epidemiological week


## Meningococcal Disease



**Figure D1**, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1**, Summary information for Meningococcal Disease (Serogroup unspecified and Invasive all serogroups)

Indicator	New Mexico
Report period	1952-2009
Total weeks	2,106
Total cases	655
Max. cases per year	66
Year (max)	1953
Max. cases per week	44
Week (max)	1952, wk 53
Average cases per year	
<i>before 1980</i>	12
<i>95%CI</i>	(6-18)
<i>after 1980</i>	11
<i>95%CI</i>	(8-14)
Average cases per week	
<i>before 1980</i>	0
<i>95%CI</i>	(0-0)
<i>after 1980</i>	0
<i>95%CI</i>	(0-0)



**Figure D2**, Number of cases reported for Meningococcal Disease per epidemiological week


## Mumps



**Figure D1**, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1, Summary information for Mumps**

Indicator	Albuquerque
Report period	1924-1932
Total weeks	423
Total cases	784
Max. cases per year	279
Year (max)	1927
Max. cases per week	33
Week (max)	1927, wk 13
Average cases per year	
<i>before 1980</i>	87
95%CI	(15-159)
Average cases per week	
<i>before 1980</i>	2
95%CI	(2-2)



**Figure D2**, Number of cases reported for Mumps per epidemiological week


## Pneumonia



**Figure D1**, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1**, Summary information for Pneumonia (Unspecified)

Indicator	New Mexico
Report period	1948-1951
Total weeks	166
Total cases	2,191
Max. cases per year	776
Year (max)	1951
Max. cases per week	65
Week (max)	1951, wk 12
Average cases per year	548
95%CI	(139-957)
Average cases per week	13
95%CI	(11-15)



**Figure D2**, Number of cases reported for Pneumonia per epidemiological week


## Poliomyelitis



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1, Summary information for Poliomyelitis (Total)**

Indicator	New Mexico
Report period	1927-1970
Total weeks	1,675
Total cases	2,459
Max. cases per year	495
Year (max)	1952
Max. cases per week	56
Week (max)	1952, wk 32
Average cases per year	
<i>before 1960</i>	72
<i>95%CI</i>	(39-105)
<i>after 1960</i>	4
<i>95%CI</i>	(1-7)
Average cases per week	
<i>before 1960</i>	2
<i>95%CI</i>	(2-2)
<i>after 1960</i>	0
<i>95%CI</i>	(0-0)



**Figure D2**, Number of cases reported for Poliomyelitis per epidemiological week


## Rabies in Animals



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1, Summary information for Rabies in Animals**

Indicator	New Mexico
Report period	1948-2009
Total weeks	1,944
Total cases	1,101
Max. cases per year	76
Year (max)	1974
Max. cases per week	24
Week (max)	1965, wk 32
Average cases per year	18
95%CI	(14-22)
Average cases per week	1
95%CI	(1-1)



**Figure D2,** Number of cases reported for Rabies in Animals per epidemiological week


## Salmonellosis



**Figure D1**, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1**, Summary information for Salmonellosis (PHLIS and Unspecified)

Indicator	New Mexico
Report period	1999-2009
Total weeks	488
Total cases	2,769
Max. cases per year	389
Year (max)	2008
Max. cases per week	92
Week (max)	2008, wk 25
Average cases per year	252
95%CI	(215-289)
Average cases per week	6
95%CI	(5-7)



**Figure D2**, Number of cases reported for Salmonellosis per epidemiological week


## Scarlet Fever



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Scarlet Fever (Including streptococcal sore throat and Unspecified)

Indicator	New Mexico	Albuquerque
Report period	1927-1969	1919-1953
Total weeks	1,782	1,020
Total cases	112,731	1,664
Max. cases per year	15,063	394
Year (max)	1964	1936
Max. cases per week	781	26
Week (max)	1966, wk 11	1936, wk 09
Average cases per year	3,047	67
95%CI	(1,443-4,651)	(34-100)
Average cases per week	63	2
95%CI	(58-68)	(2-2)



**Figure D2,** Number of cases reported for Scarlet Fever per epidemiological week


## Shigellosis



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1,** Summary information for Shigellosis (Unspecified and PHLIS)


Indicator	New Mexico
Report period	1999-2009
Total weeks	497
Total cases	1,626
Max. cases per year	251
Year (max)	2005
Max. cases per week	143
Week (max)	1999, wk 33
Average cases per year	148
95%CI	(102-194)
Average cases per week	3
95%CI	(2-4)



**Figure D2,** Number of cases reported for Shigellosis per epidemiological week

Preliminary data, not for publication or official use


# Streptococcal Disease, Invasive Group A



**Figure D1**, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1**, Summary information for Streptococcal Disease, Invasive Group A

<b>Indicator</b>	<b>New Mexico</b>
Report period	2002-2009
Total weeks	374
Total cases	574
Max. cases per year	100
Year (max)	2003
Max. cases per week	13
Week (max)	2006, wk 27
Average cases per year	72
95%CI	(54-90)
Average cases per week	2
95%CI	(2-2)



**Figure D2**, Number of cases reported for Group A Streptococcal Disease per epidemiological week


## Streptococcal Sore Throat



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1, Summary information for Streptococcal Sore Throat**

Indicator	New Mexico
Report period	1960-1961
Total weeks	99
Total cases	27,936
Max. cases per year	15,702
Year (max)	1961
Max. cases per week	802
Week (max)	1961, wk 08
Average cases per year	13,968
95%CI	(-8,065-36,001)
Average cases per week	282
95%CI	(257-307)



**Figure D2,** Number of cases reported for Streptococcal Sore Throat per epidemiological week


## Syphilis



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1,** Summary information for Syphilis (Primary and secondary and Civilian primary and secondary)

Indicator	New Mexico
Report period	1972-2009
Total weeks	1,623
Total cases	2,825
Max. cases per year	273
Year (max)	1976
Max. cases per week	90
Week (max)	1977, wk 42
Average cases per year	74
95%CI	(53-95)
Average cases per week	2
95%CI	(2-2)



**Figure D2,** Number of cases reported for Syphilis per epidemiological week


## Tuberculosis



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1,** Summary information for Tuberculosis (Unspecified)

Indicator	New Mexico
Report period	1975-2005
Total weeks	1,400
Total cases	2,855
Max. cases per year	231
Year (max)	1991
Max. cases per week	143
Week (max)	1991, wk 29
Average cases per year	92
95%CI	(73-111)
Average cases per week	2
95%CI	(2-2)



**Figure D2,** Number of cases reported for Tuberculosis per epidemiological week


## Typhoid Fever



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1,** Summary information for Typhoid Fever (Unspecified and Including paratyphoid fever)

Indicator	New Mexico	Albuquerque
Report period	1927-2003	1918-1953
Total weeks	2,506	949
Total cases	4,880	428
Max. cases per year	431	51
Year (max)	1935	1922
Max. cases per week	35	20
Week (max)	1935, wk 42	1922, wk 38
Average cases per year		
<i>before 1950</i>	182	18
95%CI	(129-235)	(13-23)
<i>after 1950</i>	12	3
95%CI	(7-17)	(3-3)
Average cases per week		
<i>before 1950</i>	4	0
95%CI	(4-4)	(0-0)
<i>after 1950</i>	0	1
95%CI	(0-0)	(0-2)



**Figure D2,** Number of cases reported for Typhoid Fever per epidemiological week


## Typhus Fever



**Figure D1**, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.


**Table D1**, Summary information for Typhus Fever (Unspecified and Endemic)

Indicator	New Mexico
Report period	1942-1947
Total weeks	197
Total cases	7
Max. cases per year	3
Year (max)	1946
Max. cases per week	1
Week (max)	1942, wk 13
Average cases per year	1
95%CI	(0-2)
Average cases per week	0
95%CI	(0-0)



**Figure D2**, Number of cases reported for Typhus Fever per epidemiological week


## Whooping Cough



**Figure D1,** Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

**Table D1,** Summary information for Whooping Cough

Indicator	New Mexico	Albuquerque
Report period	1937-2009	1924-1953
Total weeks	2,233	893
Total cases	15,195	1,745
Max. cases per year	1,493	501
Year (max)	1940	1929
Max. cases per week	150	57
Week (max)	2005, wk 50	1929, wk 11
Average cases per year		
<i>before 1960</i>	674	87
95%CI	(487-861)	(35-139)
<i>after 1960</i>	67	-
95%CI	(46-88)	-
Average cases per week		
<i>before 1960</i>	15	2
95%CI	(14-16)	(2-2)
<i>after 1960</i>	2	-
95%CI	(2-2)	-



**Figure D2,** Number of cases reported for Whooping Cough per epidemiological week

## Diseases with fragmented data over time

For a number of diseases, only fragmented data was available for New Mexico. This was due to the inclusion of these diseases in the weekly surveillance system for short periods or widely dispersed periods of time. Due to these wide gaps or limited data, data for these diseases were not presented in separate sections but in the table below. Work will continue to include more data in the Tycho database and complete data sets for each disease where possible. Collaborations with local, state and federal public health agencies will be made to explore data availability and opportunities to include these in the Tycho database.

**Table 2**, Summary information on the occurrence of diseases with fragmented data in New Mexico

Disease	Report type	Report period	Number of reports	Total cases
Anthrax	State	1942-1945	143	0
Botulism	State	1952-1952	1	1
Brucellosis [undulant fever]	City	1953-1953	1	1
Brucellosis [undulant fever]	State	1943-1977	145	314
Dengue	City	1924-1924	2	0
Dysentery	State	1942-1948	277	388
Encephalitis	City	1953-1953	1	1
Encephalitis	State	1942-1993	635	227
Influenza	City	1920-1941	334	1089
Influenza	State	1919-1951	734	21928
Leprosy	State	1942-1989	168	0
Lyme disease	State	1992-2009	382	25
Malaria	State	1952-2008	1353	187
Pellagra	City	1924-1932	62	11
Psittacosis	State	1958-1959	2	2
Rocky mountain spotted fever	City	1952-1952	1	1
Rocky mountain spotted fever	State	1942-2009	674	57
Rubella	State	1968-1999	781	1492
Tetanus	State	1962-1967	54	1
Toxic shock syndrome	State	1984-1993	200	29
Trichiniasis	City	1953-1953	1	1
Tularemia	State	1942-1994	855	62
West nile disease	State	2003-2009	143	153

Preliminary data, not for publication or official use

## **Project Tycho**

This report provides preliminary data for the state of New Mexico available in the Tycho database. This database is currently being beta tested and these data cannot be used for publication or other official use at this time. An open access release to the general public is planned for later in 2011.

Please visit the Tycho website for more information and to query the database at: [www.tycho.pitt.edu](http://www.tycho.pitt.edu). For further information regarding the Tycho project, contact Dr. Wilbert van Panhuis at the University of Pittsburgh Graduate School of Public Health.

Dr. Wilbert van Panhuis, MD PdD  
Graduate School of Public Health  
University of Pittsburgh  
130 DeSoto Street  
704 Parran Hall  
Pittsburgh, 15261 PA  
Tel: 412-624-7693  
Email: wav10@pitt.edu