

Project Tycho

Preliminary data for the state of Wyoming

Tycho database beta test version

The data presented in this report are of preliminary nature and should not be used for publication or other types of official use

June 2011

University of Pittsburgh Graduate School of Public Health



University of Pittsburgh



Table of Content

Acknowledgements	page 3
Contact information	page 4
Introduction	page 5
Vision and goal	page 7
Collaboration	page 8
Summary of methods	page 9
Overview of data available for the state of Wyoming	page 12
Disease specific data for Wyoming	page 17
- AIDS	page 17
- Chlamydia	page 18
- Cryptosporidiosis	page 19
- Diphtheria	page 20
- Escherichia Coli	page 22
- Giardiasis	page 23
- Gonorrhea	page 24
- Haemophilus Influenzae	page 25
- Hepatitis	page 26
- Leprosy	page 28
- Listeriosis	page 29
- Lyme disease	page 30
- Measles	page 31
- Meningitis	page 33
- Mumps	page 34
- Pneumonia	page 36
- Poliomyelitis	page 37
- Salmonellosis	page 39
- Scarlet fever	page 40
- Shigellosis	page 41
- Streptococcal disease, invasive group A	page 42
- Streptococcal sore throat	page 43
- Syphilis	page 44
- Tuberculosis	page 45
- Typhoid fever	page 46
- Typhus fever	page 48
Diseases with fragmented data	page 49

Acknowledgements

The development of this project would not have been possible without data collected every week by staff from city-, county-, state- and federal public health departments. We would like to acknowledge their invaluable contribution to the overall body of surveillance data that were digitized during this project. Particularly, we acknowledge staff from the Council of State and Territorial Epidemiologists (CSTE) and the US Centers for Disease Control for their overall efforts in disease surveillance and their particular input in this project.

We also acknowledge the help of many contributors that have contributed to the digitization and processing of weekly surveillance data included in this database. From the company Digital Divide Data: Linda Thomas, Eric Gold, Heak Hok, Villa Kem and Hetel Patel. From the University of Pittsburgh Graduate School of Public Health: Steve Wisniewsky, Heather Eng, Sharon Crow, David Galloway and Jia Xu. From the University of Pittsburgh School of Information Sciences: Vladimir Zadorozhny, Ying-Feng Hsu, David Paul Koenig, Fatimah Ridwan and Divyasheel Sharma. From the Johns Hopkins Bloomberg School of Public Health: Derek Cummings. From the Pittsburgh Supercomputer Center: David Graham and Andrew Adams.

This project was made possible with funding from the Bill & Melinda Gates Foundation through the Vaccine Modeling Initiative (grant ID 49276).

The Tycho development team at the University of Pittsburgh Graduate School of Public Health:

Willem G. van Panhuis

John Grefenstette

Shawn T. Brown

Su Yon Jung

Nian Shong Chok

Anne Cross

Donald S. Burke

Contact information

The Tycho database is currently being tested and a login account will be required to use the database and website (www.tycho.pitt.edu). All State Epidemiologists, CDC partners, other federal agencies and our research collaborators will be provided with user accounts. User accounts can also be provided upon specific request. Note that the quality of the data during the testing phase cannot be guaranteed to be sufficient for publication or official use.

Point of contact for the Tycho project:

Dr. Willem G. van Panhuis, MD PhD
Graduate School of Public Health
University of Pittsburgh
130 DeSoto Street
704 Parran Hall
Pittsburgh PA, 15261
Email: wav10@pitt.edu
Tel: 412-624-7693

Introduction

Project Tycho started in 2009 as part of the Vaccine Modeling Initiative (VMI) at the University of Pittsburgh Graduate School of Public Health, funded by the Bill & Melinda Gates foundation. This project aims to contribute to the availability of public health data for research and policy making. The vision for this project as described in the next section is a central access point for public health data of a detailed spatial and temporal scale for all countries. Currently, the Tycho database contains 10 million records that each represent a weekly report from a location for a specific disease. These records were extracted from weekly US Nationally Notifiable Disease Surveillance Reports between 1888 and 2009 (6300 reports) using double data entry (200 million keystrokes). The current database includes a total of 90 million reported cases and 4 million reported deaths due to notifiable diseases in the United States for the last 122 years.

Project **Tycho** is named after the Danish nobleman Tycho Brahe (1546 – 1601), who meticulously collected astronomical data. After Brahe's death, his assistant Kepler used these data to develop area laws of planetary motion.

Background

"No health department, State or local, can effectively prevent or control diseases without knowledge of when, where and under what conditions cases are occurring"

(Quote stated above weekly surveillance reports published in the Public Health Reports, 1888-1951)

This quote has accompanied weekly reports on the prevalence of notifiable diseases in the United States as long as these were published in the Public Health Reports, emphasizing the importance of data for disease control policy making. In the current day and age, this statement has lost none of its significance. On the contrary, as infectious diseases continue to pose challenges to the global public health system in an increasingly interconnected global society. The public health system has responded to this challenge by evermore sophisticated disease surveillance systems. The availability of internet and information technology has facilitated collection of detailed data on infectious disease incidence, the spread of pathogens, disease determinants, health behavior, etc. New technologies such as remote sensing and mobile reporting systems have advanced measurement of these factors in both high and low resource settings. The enormous wealth of public health information that is currently being accrued also poses its own challenges for data storage, management, preservation and dissemination. Public health agencies will need to allocate resources and develop capacity to specifically address these issues.

Although the importance of public health data collection has been widely recognized, the dissemination of data has been neglected for many reasons. The above quote does not only imply data collection but also its use and dissemination as knowledge on disease occurrence can only be derived from data after processing and analysis. Not only has disease surveillance become more sophisticated, analytical tools have changed as well and now include computational models to assess disease transmission and to evaluate disease control options. These methods have introduced the opportunity for data intensive studies of fine spatial-temporal patterns using dynamic models. Disease surveillance data at such scale are often not available for analysis for multiple reasons. First, surveillance data are not collected for research purposes but for disease monitoring and planning. Secondly, public health staff often lack time or capacity to prepare these data for analysis. Third, surveillance methodology often lacks transparency and standardization, reducing options for scientific analyses. Fourth, disease surveillance data are often

kept confidential and ownership or data use requirements are often unclear. Finally, archiving and preservation of surveillance data are often not a priority and disaggregated records may get lost, losing the opportunity to ever use these data for analysis.

The Tycho project aims at overcoming some of these challenges by providing open access to large public health datasets of great spatial-temporal detail. Open access to these data will provide great opportunities for scientific analysis and better informed policy making on disease control.

The Tycho database and website

As described in the methods section, a large heterogeneity in reporting practice was observed over the past 122 years of weekly disease surveillance and substantial efforts were made to clean and standardize these data. The current online database includes all cleaned and standardized data in a format that allows comparisons across time and space. It does not include parts of the data that require additional cleaning and research such as classification of disease subcategories.

Each record in the database provides a reported number of cases or deaths due to a specific disease for a specific location, for a certain week. The database can be searched online (www.tycho.pitt.edu) by selecting a dataset, an aggregation method and an output format. Data can be viewed in tables, graphics and maps and these can be downloaded for further use.

Data availability

The data provided by the Tycho database reflect the weekly US Nationally Notifiable Disease Surveillance System between 1888 and 2009. It includes all diseases that were reported as part of this system, but it does not yet include records from other surveillance systems such as the annual system or disease specific subsystem (such as the influenza surveillance system).

The available data also reflects changes in reporting practices over time. This explains why city or state level reports are only available for a certain time period (1888-1953 for cities and 1927-2009 for states) and why morbidity and mortality reports are available for different time periods. We are currently working on inclusion of additional data as described in the section on collaboration.

Testing and release of the Tycho database

The Tycho database is currently in a beta testing phase that will start during the 2011 annual meeting of the Council of State and Territorial Epidemiologists (CSTE) in Pittsburgh June 12-16. This phase will end with a release to the general public in the fall of 2011. During the testing phase, invited users will be provided with login accounts that will enable full use of all features of the website and database. Invited users will include all state epidemiologists, partners in the CDC and other federal agencies as well as research collaborators. During this phase, data cleaning and standardization will continue. The quality of the data will not be sufficient for publication or official use during this phase.

Future developments

After optimization of the database and website during the testing phase, the first version of the Tycho database will be released to the general public. After that, efforts to clean and standardize the data will continue and more data will be released in subsequent versions. In addition, ongoing and new collaborations for inclusion of new data will allow gradual expansion of the dataset over time and increased opportunities for data driven analysis and policy making.

Vision and goal

The vision for project Tycho is based on the value of open access to detailed, disaggregated public health data for scientific analysis and policy making. Starting with 122 years of weekly US surveillance data, it is our hope that this example will convince public health authorities worldwide of the value of this resource and will stimulate similar contributions from other countries.

In the past, data sharing and archiving have not been a priority in public health or among health sciences in general. Continuous and diligent public health data collection has led to a large pool of disaggregated data scattered over the world including the internet, ranging from paper archives to basements of local health departments. The need for open access to a central repository of public health data has been recognized and both benefits and challenges of open access to data in general have been well described. Open access to data could lead to increased accountability, transparency, innovation, collaboration, cost-effectiveness, replication of results, the development of new methods and insights, and a narrower data access gap between low- and higher income countries. Ultimately, this would result in the advancement of science and technology to the benefit of all.

Challenges to open access include inconsistent formats, lack of annotation or metadata, lack of information technology (IT) capacity, a lack of incentive, and cultural norms around data ownership. Open access to public health data in particular is confronted by decentralized archiving of disease reports, lack of time and capacity for data management, degraded physical format of records and archives, and governmental level legal constraints. Multiple principles and requirements for a central open access repository for public health data have been described and include (1) capacity building for data management in low- and middle income countries, (2) feedback loops to data contributors, (3) sustainability, (4) common standards, (5) interoperability, and (6) user-friendliness.

The ultimate goal of the Tycho project is to provide a central global public health data access point. Historical as well as current public health data are of great value if archived and accessible for research and analysis. Open access will enable the use of analytical capacity from around the globe which will lead to new discoveries of disease patterns and control policies. Furthermore, a central data access point will facilitate data archiving and preservation into the future which will be an increasing need in a data rich public health environment.

Collaboration

The application of the surveillance data from the Tycho database as well as the inclusion of new data will require a wide range of collaborations with public health offices in the US and abroad. The Tycho project has currently made the first step of digitizing 122 years of weekly US surveillance reports that had already been published. The next step will be the completion of the existing dataset as well as the inclusion of new data from unpublished sources.

This report provides an overview of data available from the Tycho database for the state of Wyoming. This includes all data that has ever been published at state or city level for Wyoming in the weekly US Nationally Notifiable Disease Surveillance System. As you will find, the data availability will vary greatly over time, between city and state level reports, between diseases and between morbidity vs. mortality reports.

The Tycho development team would look forward to work with state health departments in the US to:

1. Better understand the current Tycho data for each state (eg. by collecting historic documentation)
2. Provide better usability and applications of current Tycho data for each state by continued joint development of website and database features.
3. Collect and/or digitize official, confirmed data to validate the current (preliminary) weekly data.
4. For each state, collect and/or digitize new data that has not yet been included. For example all city level reports discontinued in 1953 and completion of city level data until 2009 would greatly increase opportunities for analysis of disease patterns and trends.
5. Provide support to state health offices to manage, preserve and provide access to public health data.

Summary of methods

Detailed documentation on the methods used for the Tycho database has been provided on the website (www.tycho.pitt.edu). This section describes these methods in short.

Data collection

Weekly reports that contain tables on the occurrence of nationally notifiable diseases have been published since 1888 by public health authorities at the Federal level in various journals. Table 1 provides the list of publications and the responsible agency since 1888. All weekly nationally notifiable disease reports between 1888 and 1951 could be retrieved from the PubMed Central repository of the National Library of Medicine ¹. For 1995 to 2009, these reports could be retrieved from the MMWR digital archive on the CDC website ². Most weekly reports between 1952 and 1995 could be retrieved from the HathiTrust Digital Library ^{3 4}, but many could not be found and had to be copied from hard copies of MMWR issues in the University of Pittsburgh library.

Table 1, publications and responsible Federal agencies for nationally notifiable disease reports

Time period	Publication title	Responsible federal agency
1888-1889	Weekly Abstract of Sanitary Reports	US Marine Hospital Service
1890-1895	Abstract of Sanitary Reports	US Marine Hospital Service
1896-1901	Public Health Reports	US Marine Hospital Service
1902-1911	Public Health Reports	US Public Health and Marine Hospital Service
1912-1951	Public Health Reports	US Public Health Service
1952-1960	Morbidity and Mortality Weekly Report	National Office of Vital Statistics, US Public Health Service
1961-1969	Morbidity and Mortality Weekly Report	Communicable Disease Center
1970-1991	Morbidity and Mortality Weekly Report	Center for Disease Control
1992-2009	Morbidity and Mortality Weekly Report	Centers for Disease Control and Prevention

Inclusion criteria

Weekly reports of each year were reviewed systematically to assess the diseases reported. We included all tables that provided disease specific information by week for US cities, townships, counties or states. Tables that provided summary or aggregated information by month, year or at the national level were not included. Similarly, tables that did not contain disease specific information (such as all cause mortality) were not included.

Data entry

Weekly reports were downloaded or scanned as PDF files and selected tables with notifiable disease reports were entered into computer spreadsheets in a highly standardized fashion using double data entry. During the second round of data entry, operators could not see what had been entered in the first round and could not continue if the system detected a discrepancy between the second and first entry

¹ <http://www.pubmedcentral.nih.gov/tocrender.fcgi?journal=333&action=archive>

² <http://www.cdc.gov/mmwr>

³ <http://catalog.hathitrust.org/Record/003910026>

⁴ <http://catalog.hathitrust.org/Record/003843660>

for a specific value. Such discrepancies could only be resolved by checking the PDF file and try again or discussion with the group leader and verification of a value in the source documents.

Quality control for data entry

The accuracy of data entry was checked at various levels. First, completeness of data was verified by comparing the content of entered data with PDF sources files. Secondly, accuracy of data entry was verified by multiple rounds of comparing random samples of entered files with PDF source files. Thirdly, data formatting was verified by various checks to ensure appropriate formatting for data loading.

Data loading and standardization

All data was entered in Excel spreadsheets and various components of these spreadsheets were loaded in data files. Table titles, column headers, place names and reported numbers were loaded in separate files. These files were used to extract information on each reported number, including:

1. the disease reported
2. the disease subcategory reported
3. cases or deaths reported
4. the reporting location (name, state and type of location)
5. the time period for which a number was reported
6. the date of publication of the original weekly report associated with a reported number

Integration

All reported numbers and extracted information was integrated in one database with one record per reported number and associated information.

Post-processing quality control

After integration of all data in one database, checks were performed to detect duplicate reports and data inconsistencies. Duplicate records were removed and inconsistencies resolved by verification with original PDF source files.

Data filtering

The digitized version of all historical weekly US nationally notifiable disease surveillance records is a very heterogeneous dataset. It took a substantial number of data processing protocols to standardize time and space variables as well as disease names. All standardized records have been separated from non-standard records by a filter. All remaining heterogeneity in non-standard records is inherent to the surveillance system and can only be standardized after further analysis will have been completed (eg. remaining heterogeneity in reporting periods for reports before 1953 and standardization of disease subcategories). The largest proportion of data (>4 million records) has been standardized however and has been made available in the current testing version.

Data visualization

Figures in this report were made with the R system, version 2.9.2 and the maps were generated by the GAIA platform developed at the University of Pittsburgh Graduate School of Public Health in collaboration with the Pittsburgh Supercomputing Center (PSC). See <http://midas-pitt.psc.edu/gaia> for more information.




Figure 1, Schematic of the Tycho data architecture. Access to a consistent, standardized subset of the digitized US weekly surveillance reports will be provided for beta testing through a website that will allow querying and downloading of data. Data from the development database will be pushed through a filter to feed the Tycho database.

The current Tycho database will provide the reported number of cases or deaths reported by city or state health authorities to the federal health agency for all weeks between 1888 and 2009. Currently, only standardized, consistent data are being released for testing, according to the following criteria:

1. Only weekly reports are provided (reports for other periods such as 10 days, 2 weeks, 1 month, 1 year) are currently not included due to lack of comparability with the weekly reports.
2. Only reports that were published in the same year as the reporting period were included (this excludes updates or comparisons that were published more than a year after the original reporting period)
3. Only numbers were included for which information about the location, time period and disease could be extracted.
4. Only numbers or time periods for which no disease subcategories were reported were included.

Each of the steps described above have been described in detailed protocols that are posted in the documentation part of the Tycho website (www.tycho.pitt.edu).

Website

The Tycho website features simple and advanced searches of the data, visualizations and downloading of machine readable files. A dataset can be selected by specifying: 1) the disease of interest, 2) the location(s) of interest and 3) the time period. In a second step, the type of aggregation and length of increments can be specified and in a third step, the type of output (table, graphic or map) can be selected. After output has been generated, data can be downloaded.

As mentioned above, data availability varies largely by disease, time period and location. We aimed to restrict selection options by data availability to avoid "no results available."

Access to the most of the website will be password protected during the testing phase. For invited users after login, full functionality of the database and website will be available. During this phase however, the data quality will not be sufficient for publication or official use.

Overview of data available for the state of Wyoming

In this section, an overview of data available for the state of Wyoming will be provided. As described earlier, data availability depends entirely on historical reporting practices in the weekly US Nationally Notifiable Disease Surveillance System between 1888 and 2009. For this report, we only included morbidity reports (cases) to improve the format and limit the size of this report. As mentioned earlier, the analysis and standardization of has not been completed yet for all records in the database and preliminary data are provided here that may not yet be available in the online database.

This section will provide a general overview of data availability. The next sections will provide disease specific data from state and city level reports of Wyoming. Data for a maximum of 3 major cities were provided depending on availability (city data was only reported until 1953). Some summary data will be provided at the end of this report, for a subset of diseases for which data was fragmented over time.

Table 1 lists the number of weekly state or city reports that are available for Wyoming per each disease and subcategory.

Table 1, Number of weekly state or city reports per disease and subcategory

Disease	City	State
Aids	-	826
Anthrax	-	143
Brucellosis [undulant fever]	-	148
Chickenpox [varicella]	-	383
Chlamydia	-	675
Coccidioidomycosis	-	178
Cryptosporidiosis	-	393
Diphtheria	37	940
Dysentery		
<i>Amebic</i>	-	212
<i>Bacillary</i>	-	203
<i>Unspecified</i>	-	197
Encephalitis		
<i>Post infectious</i>	-	34
<i>Primary [infectious] including unspecified</i>	-	506
Escherichia coli		
<i>EHEC 0157</i>	-	104
<i>EHEC non-0157</i>	-	124
<i>O157:H7 NETSS</i>	-	204
<i>O157:H7 PHLIS</i>	-	166
<i>STEC</i>	-	118
Giardiasis	-	370
Gonorrhea		
<i>Civilian</i>	-	754
<i>Unspecified</i>	-	1001

Table 1, Number of weekly state or city reports per disease and subcategory, *continued*

Disease	City	State
Haemophilus influenzae		
<i>Age <5 non-serotype B</i>	-	9
<i>Age <5 unknown serotype</i>	-	31
<i>All ages all serotypes</i>	-	426
Hepatitis		
<i>Acute type A</i>	-	249
<i>Acute type B</i>	-	248
<i>Acute type C</i>	-	28
<i>Acute type NA NB [including C]</i>	-	42
<i>All types, <20 years</i>	-	84
<i>All types, ≥20 years</i>	-	39
<i>All types, all ages</i>	-	582
<i>Type A [infectious]</i>	-	1043
<i>Type B [serum]</i>	-	686
<i>Type NA NB [including C]</i>	-	599
<i>Type unspecified</i>	-	109
Influenza	17	368
Legionellosis	-	666
Leprosy	-	155
Listeriosis	-	43
Lyme disease	-	507
Malaria	-	544
Measles		
<i>Imported</i>	-	85
<i>Indigenous</i>	-	97
<i>Unspecified</i>	53	2179
Meningitis		
<i>Aseptic</i>	-	262
<i>Meningococcus</i>	-	1127
<i>Unspecified</i>	3	5
Meningococcal disease		
<i>Invasive all serogroups</i>	-	157
<i>Invasive serogroup unknown</i>	-	35
<i>Serogroup unspecified</i>	-	1297
Mumps	-	952
Pneumonia	36	108
Poliomyelitis		
<i>Non paralytic</i>	-	40
<i>Paralytic</i>	-	146
<i>Total</i>	2	1540
Psittacosis	-	1

Table 1, Number of weekly state or city reports per disease and subcategory, *continued*

Disease	City	State
Rabies in animals	-	1526
Rocky mountain spotted fever	-	1040
Rubella	-	596
Salmonellosis		
<i>NETSS</i>	-	126
<i>PHLIS</i>	-	118
<i>Unspecified</i>	-	376
Scarlet fever		
<i>Including streptococcal sore throat</i>	-	538
<i>Unspecified</i>	81	1188
Shigellosis		
<i>NETSS</i>	-	95
<i>PHLIS</i>	-	88
<i>Unspecified</i>	-	310
Streptococcal disease, invasive group a	-	345
Streptococcal sore throat	-	73
Streptococcus pneumoniae invasive disease		
<i>Drug resistant <5 years</i>	-	69
<i>Drug resistant A</i>	-	5
<i>Drug resistant all ages</i>	-	329
<i>Non drug resistant <5 years</i>	-	32
Syphilis		
<i>Civilian primary and secondary</i>	-	615
<i>Primary and secondary</i>	-	566
Tetanus	-	73
Toxic shock syndrome	-	91
Trichiniasis	-	1
Tuberculosis [phthisis pulmonalis]		
<i>New active</i>	-	112
<i>Unspecified</i>	50	1046
Tularemia	-	999
Typhoid fever [enteric fever]		
<i>Including paratyphoid fever</i>	-	558
<i>Unspecified</i>	11	1146
Typhus fever		
<i>Endemic</i>	-	21
<i>Unspecified</i>	-	176
West nile disease		
<i>Neuroinvasive</i>	-	112
<i>Non-neuroinvasive</i>	-	96
Whooping cough [pertussis]	-	1630

The diseases included in the weekly US Nationally Notifiable Disease Surveillance System varied largely over time and reflected the historical social-political priorities of each time period. Note that the diseases in the weekly system were a subset of all diseases included in the annual Notifiable Disease Surveillance System (that was not entered as part of this project).

The number of locations reporting diseases also varied over time depending on the type of reports submitted. Figure 1 provides an overview of the weeks for which a report was available per disease at the state or city level. Colors indicate the total number of reporting locations.




Figure 1, The number of locations (in color code) reporting on each disease per week between 1888 and 2009

The map in figure 2 (next page) illustrates the geographical distribution of the available city data. It displays the number of weekly reports for any disease available per location that was included (and for which coordinates could be derived).

Figure 3 below provides an overview of the number of diseases for which a weekly report was included at the state level and for individual cities. All cities for which at least 100 weekly reports were available have been listed. For most states, it shows a major drop in the number of cities after 1925 and after 1953, city reports were discontinued except for New York City, which became its own reporting jurisdiction.




Figure 2, This map displays the number of weekly reports available for any disease per city (note that city reports were only available until 1953)




Figure 3, The number of diseases (in color code) that were reported for the state and city level (state on top row) for each week between 1888 and 2009

Disease specific data for Wyoming

AIDS




Figure D1, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

Table D1, Summary information for AIDS

Indicator	Wyoming
Report period	1984-2005
Total weeks	826
Total cases	292
Max. cases per year	109
Year (max)	2004
Max. cases per week	94
Week (max)	2004, wk 22
Average cases per year	14
95%CI	(3-25)
Average cases per week	0
95%CI	(0-0)




Figure D2, Number of cases reported for AIDS per epidemiological week

Chlamydia




Figure D1, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

Table D1, Summary information for Chlamydia

Indicator	Wyoming
Report period	1996-2009
Total weeks	675
Total cases	14,431
Max. cases per year	3,685
Year (max)	2004
Max. cases per week	2,705
Week (max)	2004, wk 22
Average cases per year	1,031
95%CI	(561-1,501)
Average cases per week	21
95%CI	(13-29)




Figure D2, Number of cases reported for Chlamydia per epidemiological week

Cryptosporidiosis




Figure D1, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

Table D1, Summary information for Cryptosporidiosis

Indicator	Wyoming
Report period	1999-2009
Total weeks	393
Total cases	179
Max. cases per year	54
Year (max)	2007
Max. cases per week	22
Week (max)	2004, wk 22
Average cases per year	16
95%CI	(4-28)
Average cases per week	0
95%CI	(0-0)




Figure D2, Number of cases reported for Cryptosporidiosis per epidemiological week

Diphtheria




Figure D1, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

Table D1, Summary information for Diphtheria

Indicator	Wyoming
Report period	1927-1962
Total weeks	940
Total cases	846
Max. cases per year	111
Year (max)	1928
Max. cases per week	16
Week (max)	1962, wk 43
Average cases per year	
<i>before 1940</i>	40
<i>95%CI</i>	(25-55)
<i>after 1940</i>	15
<i>95%CI</i>	(8-22)
Average cases per week	
<i>before 1940</i>	2
<i>95%CI</i>	(2-2)
<i>after 1940</i>	0
<i>95%CI</i>	(0-0)




Figure D2, Number of cases reported for Diphtheria per epidemiological week

Escherichia Coli




Figure D1, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

Table D1, Summary information for Escherichia Coli (O157:H7 PHLIS, EHEC 0157, and STEC)

Indicator	Wyoming
Report period	1996-2009
Total weeks	388
Total cases	96
Max. cases per year	19
Year (max)	2006
Max. cases per week	9
Week (max)	1999, wk 43
Average cases per year	7
95%CI	(4-10)
Average cases per week	0
95%CI	(0-0)




Figure D2, Number of cases reported for Escherichia Coli per epidemiological week

Giardiasis




Figure D1, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

Table D1, Summary information for Giardiasis

Indicator	Wyoming
Report period	2002-2009
Total weeks	370
Total cases	1,125
Max. cases per year	907
Year (max)	2006
Max. cases per week	880
Week (max)	2006, wk 35
Average cases per year	141
95%CI	(-118-400)
Average cases per week	3
95%CI	(-2-8)




Figure D2, Number of cases reported for Giardiasis per epidemiological week

Gonorrhea




Figure D1, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

Table D1, Summary information for Gonorrhea (Unspecified and Civilian)

Indicator	Wyoming
Report period	1972-2009
Total weeks	1,755
Total cases	13,316
Max. cases per year	1,067
Year (max)	1980
Max. cases per week	82
Week (max)	2006, wk 36
Average cases per year	350
95%CI	(231-469)
Average cases per week	8
95%CI	(8-8)




Figure D2, Number of cases reported for Gonorrhea per epidemiological week

Haemophilus Influenzae




Figure D1. Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

Table D1, Summary information for Haemophilus Influenzae (All ages all serotypes)

Indicator	Wyoming
Report period	1996-2009
Total weeks	426
Total cases	97
Max. cases per year	34
Year (max)	1996
Max. cases per week	23
Week (max)	2006, wk 35
Average cases per year	7
95%CI	(1-13)
Average cases per week	0
95%CI	(0-0)




Figure D2, Number of cases reported for Haemophilus Influenzae per epidemiological week

Hepatitis




Figure D1, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

Table D1, Summary information for Hepatitis (Type A [Infectious], Acute type A, Type B [Serum], and Acute type B)

Indicator	Wyoming
Report period	1966-2009
Total weeks	1,436
Total cases	1,540
Max. cases per year	119
Year (max)	1970
Max. cases per week	36
Week (max)	1970, wk 44
Average cases per year	
<i>before 1990</i>	45
<i>95%CI</i>	(34-56)
<i>after 1990</i>	22
<i>95%CI</i>	(11-33)
Average cases per week	
<i>before 1990</i>	2
<i>95%CI</i>	(2-2)
<i>after 1990</i>	1
<i>95%CI</i>	(1-1)




Figure D2, Number of cases reported for Hepatitis per epidemiological week

Leprosy




Figure D1, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

Table D1, Summary information for Leprosy

Indicator	Wyoming
Report period	1942-1945
Total weeks	155
Total cases	0

Listeriosis



Figure D1, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

Table D1, Summary information for Listeriosis

Indicator	Wyoming
Report period	2000-2001
Total weeks	43
Total cases	1
Max. cases per year	1
Year (max)	2001

Lyme Disease




Figure D1, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

Table D1, Summary information for Lyme Disease

Indicator	Wyoming
Report period	1991-2009
Total weeks	507
Total cases	52
Max. cases per year	8
Year (max)	1993
Max. cases per week	8
Week (max)	2000, wk 35
Average cases per year	3
95%CI	(2-4)
Average cases per week	0
95%CI	(0-0)




Figure D2, Number of cases reported for Lyme Disease per epidemiological week

Measles




Figure D1, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

Table D1, Summary information for Measles (Unspecified)

Indicator	Wyoming
Report period	1927-1996
Total weeks	2,179
Total cases	37,115
Max. cases per year	2,883
Year (max)	1943
Max. cases per week	308
Week (max)	1951, wk 19
Average cases per year	
<i>before 1970</i>	838
<i>95%CI</i>	(619-1,057)
<i>after 1970</i>	20
<i>95%CI</i>	(2-38)
Average cases per week	
<i>before 1970</i>	20
<i>95%CI</i>	(18-22)
<i>after 1970</i>	1
<i>95%CI</i>	(1-1)




Figure D2, Number of cases reported for Measles per epidemiological week

Meningitis




Figure D1, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

Table D1, Summary information for Meningitis (Meningococcus)

Indicator	Wyoming
Report period	1927-1964
Total weeks	1,127
Total cases	339
Max. cases per year	53
Year (max)	1928
Max. cases per week	10
Week (max)	1928, wk 01
Average cases per year	13
95%CI	(8-18)
Average cases per week	0
95%CI	(0-0)




Figure D2, Number of cases reported for Meningitis per epidemiological week

Mumps




Figure D1, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

Table D1, Summary information for Mumps

Indicator	Wyoming
Report period	1968-2001
Total weeks	952
Total cases	1,728
Max. cases per year	436
Year (max)	1973
Max. cases per week	150
Week (max)	1968, wk 05
Average cases per year	
<i>before 1980</i>	153
<i>95%CI</i>	(19-287)
<i>after 1980</i>	3
<i>95%CI</i>	(2-4)
Average cases per week	
<i>before 1980</i>	4
<i>95%CI</i>	(3-5)
<i>after 1980</i>	0
<i>95%CI</i>	(0-0)




Figure D2, Number of cases reported for Mumps per epidemiological week

Pneumonia




Figure D1, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

Table D1, Summary information for Pneumonia (Unspecified)

Indicator	Wyoming
Report period	1948-1951
Total weeks	108
Total cases	408
Max. cases per year	217
Year (max)	1949
Max. cases per week	20
Week (max)	1949, wk 17
Average cases per year	102
95%CI	(-27-231)
Average cases per week	4
95%CI	(3-5)




Figure D2, Number of cases reported for Pneumonia per epidemiological week

Poliomyelitis




Figure D1, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

Table D1, Summary information for Poliomyelitis (Total)

Indicator	Wyoming
Report period	1927-1970
Total weeks	1,540
Total cases	1,474
Max. cases per year	273
Year (max)	1954
Max. cases per week	26
Week (max)	1951, wk 37
Average cases per year	
<i>before 1960</i>	43
<i>95%CI</i>	(22-64)
<i>after 1960</i>	2
<i>95%CI</i>	(1-3)
Average cases per week	
<i>before 1960</i>	1
<i>95%CI</i>	(1-1)
<i>after 1960</i>	0
<i>95%CI</i>	(0-0)




Figure D2, Number of cases reported for Poliomyelitis per epidemiological week

Salmonellosis




Figure D1, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

Table D1, Summary information for Salmonellosis (PHLIS and Unspecified)

Indicator	Wyoming
Report period	1999-2009
Total weeks	494
Total cases	623
Max. cases per year	94
Year (max)	2002
Max. cases per week	30
Week (max)	2002, wk 45
Average cases per year	57
95%CI	(44-70)
Average cases per week	1
95%CI	(1-1)




Figure D2, Number of cases reported for Salmonellosis per epidemiological week

Scarlet Fever




Figure D1, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

Table D1, Summary information for Scarlet Fever (Including streptococcal sore throat and Unspecified)

Indicator	Wyoming
Report period	1927-1969
Total weeks	1,726
Total cases	51,810
Max. cases per year	11,268
Year (max)	1969
Max. cases per week	885
Week (max)	1969, wk 44
Average cases per year	1,400
95%CI	(638-2,162)
Average cases per week	30
95%CI	(27-33)




Figure D2, Number of cases reported for Scarlet Fever per epidemiological week

Shigellosis




Figure D1. Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

Table D1, Summary information for Shigellosis (Unspecified and PHLIS)

Indicator	Wyoming
Report period	1999-2009
Total weeks	398
Total cases	173
Max. cases per year	85
Year (max)	2006
Max. cases per week	21
Week (max)	1999, wk 33
Average cases per year	16
95%CI	(-1-33)
Average cases per week	0
95%CI	(0-0)





Figure D2, Number of cases reported for Shigellosis per epidemiological week



Indicator	Wyoming
Report period	2002-2009
Total weeks	345
Total cases	31
Max. cases per year	8
Year (max)	2004
Max. cases per week	3
Week (max)	2002, wk 19
Average cases per year	4
95%CI	(2-6)
Average cases per week	0
95%CI	(0-0)



Streptococcal Sore Throat



Figure D1, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

Table D1, Summary information for Streptococcal Sore Throat

Indicator	Wyoming
Report period	1960-1961
Total weeks	73
Total cases	3,685
Max. cases per year	1,866
Year (max)	1961
Max. cases per week	312
Week (max)	1960, wk 46
Average cases per year	1,842
95%CI	(1,543-2,141)
Average cases per week	50
95%CI	(38-62)




Figure D2, Number of cases reported for Streptococcal Sore Throat per epidemiological week

Syphilis




Figure D1, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

Table D1, Summary information for Syphilis (Primary and secondary and Civilian primary and secondary)

Indicator	Wyoming
Report period	1972-2009
Total weeks	1,181
Total cases	264
Max. cases per year	48
Year (max)	1972
Max. cases per week	7
Week (max)	1972, wk 13
Average cases per year	8
95%CI	(4-12)
Average cases per week	0
95%CI	(0-0)




Figure D2, Number of cases reported for Syphilis per epidemiological week

Tuberculosis




Figure D1, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

Table D1, Summary information for Tuberculosis (Unspecified)

Indicator	Wyoming
Report period	1975-2004
Total weeks	1,046
Total cases	275
Max. cases per year	33
Year (max)	1991
Max. cases per week	30
Week (max)	1991, wk 29
Average cases per year	11
95%CI	(7-15)
Average cases per week	0
95%CI	(0-0)




Figure D2, Number of cases reported for Tuberculosis per epidemiological week

Typhoid Fever




Figure D1, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

Table D1, Summary information for Typhoid Fever (Unspecified and Including paratyphoid fever)

Indicator	Wyoming
Report period	1927-1979
Total weeks	1,704
Total cases	586
Max. cases per year	66
Year (max)	1928
Max. cases per week	33
Week (max)	1953, wk 02
Average cases per year	
<i>before 1950</i>	19
<i>95%CI</i>	(13-25)
<i>after 1950</i>	5
<i>95%CI</i>	(1-9)
Average cases per week	
<i>before 1950</i>	0
<i>95%CI</i>	(0-0)
<i>after 1950</i>	0
<i>95%CI</i>	(0-0)




Figure D2, Number of cases reported for Typhoid Fever per epidemiological week

Typhus Fever



Figure D1, Weeks between 1888 and 2009 for which data on the disease and subcategories (if applicable) are available in the Tycho database. If no subcategory was reported, a subcategory of "Unspecified" was assigned.

Table D1, Summary information for Typhus Fever (Unspecified and Endemic)

Indicator	Wyoming
Report period	1942-1947
Total weeks	197
Total cases	2
Max. cases per year	2
Year (max)	1944

Diseases with fragmented data over time

For a number of diseases, only fragmented data was available for Wyoming. This was due to the inclusion of these diseases in the weekly surveillance system for short periods or widely dispersed periods of time. Due to these wide gaps or limited data, data for these diseases were not presented in separate sections but in the table below. Work will continue to include more data in the Tycho database and complete data sets for each disease where possible. Collaborations with local, state and federal public health agencies will be made to explore data availability and opportunities to include these in the Tycho database.

Table 2, Summary information on the occurrence of diseases with fragmented data in Wyoming

Disease	Report type	Report period	Number of reports	Total cases
Anthrax	State	1942-1945	143	0
Brucellosis [undulant fever]	State	1943-1970	148	67
Chickenpox [varicella]	State	1972-2008	383	1784
Dysentery	State	1942-1948	213	47
Encephalitis	State	1942-1994	506	160
Influenza	City	1920-1923	19	201
Influenza	State	1927-1951	368	19546
Legionellosis	State	1982-2009	666	54
Malaria	State	1952-2005	544	94
Meningococcal disease	State	1951-2009	1454	227
Psittacosis	State	1961-1961	1	1
Rabies in animals	State	1948-2009	1526	1372
Rocky mountain spotted fever	State	1942-2009	1040	308
Rubella	State	1966-1995	596	1381
Streptococcus pneumoniae invasive disease	State	2007-2009	69	2
Tetanus	State	1964-1977	73	2
Toxic shock syndrome	State	1983-1992	91	9
Trichiniasis	State	1953-1953	1	1
Tularemia	State	1942-1993	999	412
West nile disease	State	2003-2009	137	234
Whooping cough [pertussis]	State	1937-2009	1630	4220

Project Tycho

This report provides preliminary data for the state of Wyoming available in the Tycho database. This database is currently being beta tested and these data cannot be used for publication or other official use at this time. An open access release to the general public is planned for later in 2011.

Please visit the Tycho website for more information and to query the database at: www.tycho.pitt.edu. For further information regarding the Tycho project, contact Dr. Wilbert van Panhuis at the University of Pittsburgh Graduate School of Public Health.

Dr. Wilbert van Panhuis, MD PdD
Graduate School of Public Health
University of Pittsburgh
130 DeSoto Street
704 Parran Hall
Pittsburgh, 15261 PA
Tel: 412-624-7693
Email: wav10@pitt.edu